

Efficacy, Immunogenicity, and Safety of a 9-Valent Human Papillomavirus Vaccine: Subgroup Analysis of Participants From Asian Countries

S. M. Garland,¹ P. Pitisuttithum,² H. Y. S. Ngan,³ C.-H. Cho,⁴ C.-Y. Lee,⁵ C.-A. Chen,⁶ Y. C. Yang,⁷ T.-Y. Chu,⁸ N.-F. Twu,⁹ R. Samakoses,¹⁰ Y. Takeuchi,¹¹ T. H. Cheung,¹² S. C. Kim,¹³ L.-M. Huang,¹⁴ B.-G. Kim,¹⁵ Y.-T. Kim,¹⁶ K.-H. Kim,¹⁷ Y.-S. Song,¹⁸ S. Lalwani,¹⁹ J.-H. Kang,²⁰ M. Sakamoto,²¹ H.-S. Ryu,²² N. Bhatla,²³ H. Yoshikawa,²⁴ M. C. Ellison,²⁵ S. R. Han,²⁶ E. Moeller,²⁵ S. Murata,²⁶ M. Ritter,²⁵ M. Sawata,²⁶ C. Shields,²⁵ A. Walia,²⁵ G. Perez,²⁵ and A. Luxembourg²⁵

¹Western Pacific Regional HPV Labnet Reference Laboratory, Department of Infectious Disease and Microbiology, Royal Women's Hospital, Murdoch Children's Research Institute, Royal Children's Hospital and Department of Obstetrics and Gynaecology, University of Melbourne, Parkville, Victoria, Australia; ²Vaccine Trial Centre, Mahidol University, Salaya, Thailand; ³Department of Obstetrics and Gynaecology, the University of Hong Kong, China; ⁴Department of Obstetrics and Gynecology, Keimyung University School of Medicine, Daegu, South Korea; ⁵Department of Gynecology, Chang Gung Memorial Hospital, Chiayi Branch; ⁶Department of Obstetrics and Gynecology, National Taiwan University Hospital, Taipei; ⁷MacKay Memorial Hospital, Taipei; ⁸Tzu Chi Medical Center, Hualien, and ⁹Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan; ¹⁰Department of Pediatrics, Phramongkutklo Hospital, Bangkok, Thailand; ¹¹Fukui General Hospital, Japan; ¹²Department of Obstetric and Gynaecology, Chinese University of Hong Kong, China; ¹³Division of Gynecologic Oncology, Ewha Womans University Mokdong Hospital, School of Medicine Ewha Womans University, Seoul, South Korea; ¹⁴Department of Pediatrics, National Taiwan University Hospital, Taipei, Taiwan; ¹⁵Department of Obstetrics and Gynecology, Samsung Medical Center, Sungkyunkwan University School of Medicine; ¹⁶Department of Obstetrics and Gynecology, University of Ulsan College of Medicine, Asian Medical Center; ¹⁷Department of Pediatrics and Center for Vaccine Evaluation and Study, Ewha Womans University College of Medicine, and ¹⁸Obstetrics and Gynecology, College of Medicine, Seoul National University, Seoul, South Korea; ¹⁹Bharati Vidyapeeth Deemed University Medical College and Hospital, Pune, India; ²⁰Department of Pediatrics, Seoul St. Mary's Hospital, College of Medicine, the Catholic University of Korea, South Korea; ²¹Department of Gynaecology, Sasaki Foundation Kyoundo Hospital and Department of Obstetrics and Gynaecology, School of Medicine, the Jikei University, Tokyo, Japan; ²²Department of Obstetrics and Gynecology, School of Medicine, Ajou University, Suwon, South Korea; ²³Department of Obstetrics and Gynaecology, All India Institute of Medical Sciences, New Delhi, India; ²⁴Ibaraki Prefectural Central Hospital, Kasama, Ibaraki, Japan; ²⁵Merck & Co., Inc., Kenilworth, New Jersey; and ²⁶MSD K.K., Tokyo, Japan

Background. A 9-valent human papillomavirus-6/11/16/18/31/33/45/52/58 (9vHPV) vaccine extends coverage to 5 next most common oncogenic types (31/33/45/52/58) in cervical cancer versus quadrivalent HPV (qHPV) vaccine. We describe efficacy, immunogenicity, and safety in Asian participants (India, Hong Kong, South Korea, Japan, Taiwan, and Thailand) from 2 international studies: a randomized, double-blinded, qHPV vaccine-controlled efficacy study (young women aged 16–26 years; NCT00543543; Study 001); and an immunogenicity study (girls and boys aged 9–15 years; NCT00943722; Study 002).

Methods. Participants (N = 2519) were vaccinated at day 1 and months 2 and 6. Gynecological samples (Study 001 only) and serum were collected for HPV DNA and antibody assessments, respectively. Injection-site and systemic adverse events (AEs) were monitored. Data were analyzed by country and vaccination group.

Results. 9vHPV vaccine prevented HPV-31/33/45/52/58–related persistent infection with 90.4%–100% efficacy across included countries. At month 7, ≥97.9% of participants seroconverted for each HPV type. Injection-site AEs occurred in 77.7%–83.1% and 81.9%–87.5% of qHPV and 9vHPV vaccine recipients in Study 001, respectively, and 62.4%–85.7% of girls/boys in Study 002; most were mild to moderate.

Conclusions. The 9vHPV vaccine is efficacious, immunogenic, and well tolerated in Asian participants. Data support 9vHPV vaccination programs in Asia.

Clinical Trials Registration. NCT00543543; NCT00943722.

Keywords. 9vHPV; Asia; cervical cancer; human papillomavirus; vaccine.

Approximately 280 000 new cervical cancers and over 140 000 related deaths occur annually in Asia (based on 2012 data [1, 2]). An additional approximately 40 000 new annual cases of other human papillomavirus (HPV)-related cancers (vaginal, vulvar,

anal, penile, and oropharyngeal cancers) are estimated to affect Asian men and women [2]. Access to cervical cancer screening is highly variable in the region, and interventions for high-grade dysplasia are generally limited [1, 3–5].

Prophylactic HPV vaccination represents a unique opportunity to reduce the burden of HPV-associated disease. Bivalent HPV-16/18 and quadrivalent HPV-6/11/16/18 (qHPV) vaccines introduced over the past decade have potential to prevent the approximately 70% of cervical cancer cases attributable to HPV-16/18, both worldwide and within Asia [6]. The qHPV vaccine also protects against the approximately 90% of genital warts cases associated with HPV-6/11 [7]. Partial and inconsistent cross-protection against phylogenetically related oncogenic HPV types (31/33/45) has been observed for both vaccines in

Received 15 November 2017; editorial decision 5 March 2018; accepted 16 March 2018; published online May 15, 2018.

Correspondence: S. M. Garland, MD, The Royal Women's Hospital, Locked Bag 300, Parkville 3052, Victoria, Australia (suzanne.garland@thewomens.org.au).

The Journal of Infectious Diseases® 2018;218:95–108

© The Author(s) 2018. Published by Oxford University Press for the Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com
DOI: 10.1093/infdis/jiy133

clinical studies [8]; however, its extent, duration, and public health significance remain uncertain [8–11]. The 9-valent HPV (6/11/16/18/31/33/45/52/58) vaccine (9vHPV vaccine; Gardasil 9, Merck & Co., Inc., Kenilworth, NJ, USA) extends coverage to HPV-31/33/45/52/58, which are next most commonly associated with cervical cancer globally [6, 12, 13]. Global epidemiological data indicate that the 9vHPV vaccine has potential to prevent approximately 90% of cervical cancers and HPV-related vulvar, vaginal, and anal cancers, 70%–85% of cervical precancerous lesions, and 90% of anogenital warts [7, 14–17]. Based on HPV subtype distribution across Asia, the 9vHPV vaccine may offer even greater cervical cancer protection (approximately 92% of cases) in Asia [6, 17], possibly because of the higher relative contributions of HPV-52 and HPV-58 (see [Supplementary Material](#) for more information).

In an international phase 2b/3 trial (Study 001 [18, 19]), the 9vHPV vaccine demonstrated 97% efficacy against HPV-31/33/45/52/58-related infection and disease and elicited noninferior antibody responses to HPV-6/11/16/18 compared with the qHPV vaccine in young women 16–26 years of age. In a separate phase 3 trial (Study 002), antibody responses to HPV-6/11/16/18/31/33/45/52/58 in girls and boys aged 9–15 years were noninferior to those among young women aged 16–26 years following 9vHPV vaccination, thereby supporting the bridging of efficacy findings with 9vHPV vaccine from young women to girls and boys [20].

These 2 studies included participants from multiple Asian countries (Hong Kong, India, Japan, South Korea, Taiwan, and Thailand), highlighting the regional importance of HPV-related disease in these countries. Given the specific epidemiology of HPV in East Asia (higher prevalence of HPV-52/58; see [Supplementary Material](#) for more information), it is of interest to assess the impact of the 9vHPV vaccine on the incidence of persistent infection and cervical cytological abnormalities, disease, and medical procedures related to these 2 HPV types in the subgroup of Asian participants: differences between the Asian subgroup and the overall study population may not be apparent in the overall analyses, as Asian participants represent 12% of the population enrolled in Study 001. Also, few studies have reported the incidence of cervical cytological abnormalities related to HPV-52 and HPV-58 in populations from East Asia. Subpopulation analyses of efficacy, immunogenicity, and safety in Asian participants enrolled in Study 001 and 002 were conducted to better understand the potential impact of the 9vHPV vaccine on reducing the burden of HPV-related infection and disease in the region, and help support public health decisions regarding HPV vaccination programs in Asia.

METHODS

Study Design and Population

Study 001 (Merck & Co., Inc., Kenilworth, NJ, USA Protocol V503-001 [NCT00543543]) [18, 19, 21–23] was a randomized, double-blind, qHPV vaccine-controlled study that

evaluated 9vHPV vaccine efficacy in 14215 young women (aged 16–26 years) across 105 sites and 18 countries worldwide, including 25 sites in 5 Asian countries (Hong Kong, Japan, South Korea, Taiwan, and Thailand; N = 1717; [Supplementary Table 1](#)). All Asian participants randomized in the efficacy study were included in the subgroup analysis; those who received low or high doses of 9vHPV vaccine during the dose selection phase (69 participants from Taiwan) were not included. Data from participants from Hong Kong and Taiwan were pooled, as these individuals were considered to be part of the same ethnic group (Chinese).

Study 002 (Merck & Co., Inc., Kenilworth, NJ, USA Protocol V503-002 [NCT00943722]) [20, 24] assessed immunogenicity and safety of the 9vHPV vaccine in girls and boys (aged 9–15 years) compared with young women (aged 16–26 years) across 72 sites in 17 countries (N = 3074), including 16 sites in 4 Asian countries (India, South Korea, Taiwan, and Thailand; N = 733; [Supplementary Table 1](#)). All randomized girls and boys from these countries were included in the subgroup analysis. Data from girls and boys were pooled within each country. Because only small numbers of women from South Korea, Taiwan, and Thailand (20, 20, and 60, respectively) were enrolled in Study 002 (only 1 site enrolled young women in each of these countries) and substantial numbers of young women from these countries had participated in Study 001 (307, 531, and 465, respectively) ([Supplementary Table 1](#)), women from Study 001 were used as the comparator. Immunogenicity and safety data for Indian young women who participated in Study 002 are described, because Study 001 did not include this country. Study 002 did not include Japanese participants; a separate immunogenicity and safety study of the 9vHPV vaccine was conducted in a group of 100 Japanese girls (Study V503-008 [NCT01254643]), as reported separately [25].

Studies 001 and 002 were performed in accordance with the principles of Good Clinical Practice and were approved by the appropriate institutional review boards and regulatory agencies. All participants provided written, informed consent before study participation in accordance with local laws and regulations.

Vaccination and Follow-up

The 9vHPV vaccine (both studies) and qHPV vaccine (control arm of Study 001) were administered as a series of 3 intramuscular doses on day 1 and at months 2 and 6.

In Study 001, cervical cytology samples and cervical and external genital swabs for detection of HPV by polymerase chain reaction (PCR) assays were collected on day 1 and at months 7, 12, and every 6 months thereafter up to month 54. Participants with cervical cytological abnormalities were referred for colposcopy, based on a protocol-specified algorithm [18, 19, 22]. Tissue samples were adjudicated by a pathology panel and tested for HPV by PCR, as previously described [22, 26, 27].

Serum levels of vaccine HPV-type antibodies were assessed at day 1 and month 7 using a 9-valent competitive Luminex immunoassay (cLIA) [28]. Antibody titers for each individual HPV type were determined through competition with type-specific monoclonal antibodies, so it is not possible to directly compare assay results across HPV types.

Following each vaccination, participants recorded their oral temperature for 5 days and injection-site and systemic adverse events (AEs) for 15 days on vaccination report cards [29]. AE causality was determined by the study site investigator as previously described [29]. Serious AEs (SAEs) were prespecified as those considered life threatening by the investigator or resulted in death, significant disability or incapacity, or new or prolonged existing hospitalization, or were a congenital anomaly, cancer, or “other important medical event”.

Statistical Analyses

Efficacy was evaluated in the per-protocol efficacy (PPE) population of Study 001, defined as participants who: (1) were seronegative on day 1 and PCR negative from day 1 through month 7 for the HPV type being analyzed; (2) received all 3 doses of assigned vaccine within 1 year; and (3) had no protocol violations that could interfere with the evaluation of vaccine efficacy, as judged by the study director. Supportive analyses were conducted in the HPV type-specific naïve (HNST) population, consisting of participants who were seronegative and PCR negative on day 1 for the HPV type being analyzed, received at least 1 dose of the 9vHPV or qHPV vaccine, and had efficacy follow-up after day 1; and in the modified intention-to-treat (mITT) population, consisting of participants who received at least 1 dose of the 9vHPV or qHPV vaccine and had efficacy follow-up after day 1. Combined incidence of HPV-31/33/45/52/58-related cervical, vulvar, and vaginal disease (any grade), combined incidence of HPV-31/33/45/52/58-related cervical cytological abnormalities, and combined incidence of HPV-31/33/45/52/58-related persistent infection, as well as incidence by HPV type, are presented using incidence rates (cases per 10 000 person-years), vaccine efficacy, and 95% confidence intervals (CIs). Vaccine efficacy was calculated as $100 \times (1 - 9vHPV \text{ vaccine incidence rate} / qHPV \text{ vaccine incidence rate})$. The 95% CI of vaccine efficacy was calculated with the use of a binomial distribution-based exact method [30].

Immunogenicity was assessed in the per-protocol immunogenicity populations from each study, which included participants who: (1) were seronegative on day 1 and (for young women aged 16–26 years) PCR-negative from day 1 through month 7 for the HPV type being analyzed; (2) received all 3 vaccinations within prespecified visit intervals and had available month 7 serology results obtained within a prespecified interval; and (3) had no protocol violations that could interfere with the evaluation of the immune response to vaccine, as judged by the study director. For each HPV type, the geometric mean

titer (GMT) and 95% CIs were estimated using an analysis of variance model, with log anti-HPV as the response and vaccination group as the fixed effect. Seroconversion rates and exact 95% CIs for a binomial proportion were also calculated. All of these evaluations were post hoc analyses; therefore, no statistical hypothesis tests were performed. Nonoverlapping 95% CIs were used as indicators of differences of immune response.

Safety was assessed in all participants who received at least 1 study vaccination and had follow-up data. AEs, SAEs, and AEs leading to discontinuation were summarized by vaccination group in subgroups of participants defined by country of residence.

RESULTS

Participants

Of the 14 215 young women randomized in the efficacy study (Study 001), 1717 (12.0%) were enrolled at sites in Asia. Participant baseline characteristics are shown in Table 1. Japanese participants tended to have more lifetime sexual partners than other Asian participants. Across countries, 31.0%–47.8% of participants were HPV positive at baseline (by serology or PCR), with highest rates in Japan and Thailand (Table 1). Proportions of participants testing positive for HPV by PCR or serology are shown in Supplementary Tables 2 and 3, respectively.

Of the 2604 girls and boys enrolled in Study 002, 608 (23.3%) were enrolled at sites in Asia (423 girls; 185 boys). Participant baseline characteristics are shown in Table 2.

Efficacy

In Study 001, the 9vHPV vaccine prevented HPV-31/33/45/52/58-related persistent (≥ 6 months) infections in Asian participants with an efficacy of 95.8% (95% CI, 87.8–98.9). Efficacy ranged from 90.4% and 94.9% among participants from Japan and Thailand, respectively, to 100% among participants from Hong Kong/Taiwan and South Korea (Table 3). Efficacy was 91.3% to 100% across the HPV types. In participants who received the qHPV vaccine, HPV-52- and HPV-58-related persistent infections were most frequent in all countries. As seen in Supplementary Table 4, the 9vHPV vaccine demonstrated efficacy against persistent infection related to HPV-52 or HPV-58 among participants from Hong Kong/Taiwan, Japan, and Thailand. No cases of HPV-52/58-related persistent infection were observed in the 9vHPV vaccine group from South Korea. The vaccine also prevented HPV-31/33/45/52/58-related persistent (≥ 12 months) infections with an efficacy of 93.9% (95% CI, 81.4–98.4) in Asian participants.

No cases of cervical, vulvar, or vaginal disease (any grade) related to HPV-31/33/45/52/58 were detected in the 9vHPV vaccination group, compared with 7 cases of cervical disease (4, 2, and 1 among participants from Hong Kong/Taiwan, Japan, and Thailand, respectively) in the qHPV vaccine group (Table 3).

The 9vHPV vaccine reduced the risk of HPV-31/33/45/52/58-related cervical cytological abnormalities by 92.1% (95% CI,

Table 1. Baseline Characteristics of Asian Participants (Young Women Aged 16–26 Years) in Study 001 by Country

| Characteristics | 9vHPV Vaccine | | | | qHPV Vaccine | | | | | |
|---|--------------------------------|-----------------|-----------------------|--------------------|-----------------|--------------------------------|-----------------|-----------------------|--------------------|-----------------|
| | Hong Kong and Taiwan (N = 345) | Japan (N = 127) | South Korea (N = 154) | Thailand (N = 232) | Total (N = 858) | Hong Kong and Taiwan (N = 346) | Japan (N = 127) | South Korea (N = 153) | Thailand (N = 233) | Total (N = 859) |
| Age, y | | | | | | | | | | |
| Mean ± SD | 23.7 ± 1.9 | 22.9 ± 2.0 | 23.5 ± 1.8 | 22.4 ± 2.7 | 23.2 ± 2.2 | 23.8 ± 2.0 | 22.9 ± 2.0 | 23.5 ± 1.9 | 22.0 ± 2.6 | 23.1 ± 2.3 |
| Median | 24.0 | 23.0 | 24.0 | 23.0 | 24.0 | 24.0 | 23.0 | 24.0 | 22.0 | 23.0 |
| Range | 19–26 | 18–26 | 19–26 | 16–26 | 16–26 | 17–26 | 18–26 | 18–26 | 16–26 | 16–26 |
| Age at first sexual intercourse, y | | | | | | | | | | |
| Mean ± SD | 18.9 ± 2.5 | 17.7 ± 1.9 | 20.9 ± 2.2 | 18.0 ± 2.7 | 18.8 ± 2.7 | 19.1 ± 2.3 | 17.6 ± 1.9 | 21.2 ± 2.2 | 17.7 ± 2.2 | 18.8 ± 2.5 |
| Smoking status, No. (%) ^a | | | | | | | | | | |
| Current smoker | 28 (8.1) | 31 (24.4) | 15 (9.7) | 17 (7.3) | 91 (10.6) | 32 (9.2) | 37 (29.1) | 14 (9.2) | 14 (6.0) | 97 (11.3) |
| Former smoker | 15 (4.3) | 12 (9.4) | 7 (4.5) | 16 (6.9) | 50 (5.8) | 14 (4.0) | 12 (9.4) | 8 (5.2) | 7 (3.0) | 41 (4.8) |
| Never smoked | 302 (87.5) | 84 (66.1) | 130 (84.4) | 199 (85.8) | 715 (83.3) | 299 (86.4) | 78 (61.4) | 130 (85.0) | 212 (91.0) | 719 (83.7) |
| Unknown | 0 (0.0) | 0 (0.0) | 2 (1.3) | 0 (0.0) | 2 (0.2) | 1 (0.3) | 0 (0.0) | 1 (0.7) | 0 (0.0) | 2 (0.2) |
| Lifetime sexual partners, No. (%) ^a | | | | | | | | | | |
| 1 | 143 (41.4) | 23 (18.1) | 54 (35.1) | 121 (52.2) | 341 (39.7) | 135 (39.0) | 21 (16.5) | 60 (39.5) | 131 (56.2) | 347 (40.4) |
| 2 | 86 (24.9) | 15 (11.8) | 45 (29.2) | 73 (31.5) | 219 (25.5) | 95 (27.5) | 21 (16.5) | 29 (19.1) | 63 (27.0) | 208 (24.2) |
| 3 | 79 (22.9) | 37 (29.1) | 24 (15.6) | 28 (12.1) | 168 (19.6) | 72 (20.8) | 33 (26.0) | 31 (20.4) | 26 (11.2) | 162 (18.9) |
| 4 | 30 (8.7) | 49 (38.6) | 21 (13.6) | 10 (4.3) | 110 (12.8) | 34 (9.8) | 42 (33.1) | 18 (11.8) | 13 (5.6) | 107 (12.5) |
| >4 | 6 (1.7) | 0 (0.0) | 1 (0.6) | 0 (0.0) | 7 (0.8) | 10 (2.9) | 0 (0.0) | 3 (2.0) | 0 (0.0) | 13 (1.5) |
| Non-HPV-related cervicovaginal infections or sexually transmitted diseases, No. (%) | | | | | | | | | | |
| Any | 15 (4.3) | 7 (5.5) | 10 (6.5) | 29 (12.5) | 61 (7.1) | 21 (6.1) | 6 (4.7) | 11 (7.2) | 23 (9.9) | 61 (7.1) |
| <i>Chlamydia</i> | 12 (3.5) | 7 (5.5) | 10 (6.5) | 28 (12.1) | 57 (6.6) | 21 (6.1) | 6 (4.7) | 11 (7.2) | 22 (9.4) | 60 (7.0) |
| <i>Gonorrhea</i> | 4 (1.2) | 0 (0.0) | 0 (0.0) | 1 (0.4) | 5 (0.6) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 4 (1.7) | 4 (0.5) |
| Contraceptive use, No. (%) ^b | | | | | | | | | | |
| Barrier | 261 (75.7) | 59 (46.5) | 64 (41.6) | 60 (25.9) | 444 (51.7) | 259 (74.9) | 60 (47.6) | 58 (38.4) | 70 (30.0) | 447 (52.2) |
| Behavior | 100 (29.0) | 80 (63.0) | 82 (53.2) | 15 (6.5) | 277 (32.3) | 96 (27.7) | 73 (57.9) | 84 (55.6) | 12 (5.2) | 265 (31.0) |
| Hormonal | 41 (11.9) | 11 (8.7) | 5 (3.2) | 140 (60.3) | 197 (23.0) | 40 (11.6) | 8 (6.3) | 6 (4.0) | 143 (61.4) | 197 (23.0) |
| Day 1 composite HPV positivity, No./total No. (%) ^c | | | | | | | | | | |
| Serologic test | 88/844 (25.6) | 47/127 (37.0) | 48/151 (31.8) | 99/232 (42.7) | 282/854 (33.0) | 73/344 (21.2) | 31/127 (24.4) | 46/149 (30.9) | 94/233 (40.3) | 244/853 (28.6) |
| PCR assay | 38/331 (11.5) | 32/126 (25.4) | 23/147 (15.6) | 44/229 (19.2) | 137/833 (16.4) | 35/335 (10.4) | 25/123 (20.3) | 30/143 (21.0) | 42/228 (18.4) | 132/829 (15.9) |
| Serologic test or PCR assay | 104/336 (31.0) | 58/127 (45.7) | 58/148 (39.2) | 110/230 (47.8) | 330/841 (39.2) | 88/335 (26.3) | 46/125 (36.8) | 56/145 (38.6) | 106/231 (45.9) | 296/836 (35.4) |

Abbreviations: 9vHPV, 9-valent human papillomavirus; PCR, polymerase chain reaction; SD, standard deviation; qHPV, quadrivalent human papillomavirus.

^aThe percentages for the number of lifetime sexual partners were calculated on the basis of the number of participants for whom there were data on sexual history at enrollment.

^bParticipants may have used more than 1 contraceptive method. A participant is counted once within a category and may be counted in more than 1 category. The percentages for the numbers of participants who used contraceptives were based on the number for whom this information was available.

^cPositivity was defined as an anti-HPV titer on immunoassay of at least 30, 16, 20, 24, 10, 8, 8, and 8 for HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58, respectively. The numerator in this category represents the number of HPV-positive participants, and the denominator the total number of participants with assay results that could be evaluated.

Table 2. Baseline Characteristics of Asian Participants (Girls and Boys Aged 9–15 Years) in Study 002 by Country

| Characteristics | 9vHPV Vaccine | | | | |
|------------------------------|-----------------|-----------------------|------------------|--------------------|-----------------|
| | India (N = 200) | South Korea (N = 129) | Taiwan (N = 139) | Thailand (N = 140) | Total (N = 608) |
| Gender, No. (%) | | | | | |
| Male | 75 (37.5) | 40 (31.0) | 30 (21.6) | 40 (28.6) | 185 (30.4) |
| Female | 125 (62.5) | 89 (69.0) | 109 (78.4) | 100 (71.4) | 423 (69.6) |
| Age | | | | | |
| 9–12 y, No. (%) | 137 (68.5) | 90 (69.8) | 87 (62.6) | 81 (57.9) | 395 (65.0) |
| 13–15 y, No. (%) | 63 (31.5) | 39 (30.2) | 52 (37.4) | 59 (42.1) | 213 (35.0) |
| Mean, y ± SD | 11.5 ± 1.8 | 11.6 ± 1.8 | 11.7 ± 1.9 | 12.0 ± 1.7 | 11.7 ± 1.8 |
| Median, y | 11.0 | 12.0 | 11.0 | 12.0 | 12.0 |
| Range, y | 9–15 | 9–15 | 9–15 | 9–15 | 9–15 |
| Weight, kg | | | | | |
| Mean ± SD | 36.8 ± 9.8 | 47.2 ± 15.0 | 44.8 ± 12.6 | 42.6 ± 12.1 | 42.1 ± 12.8 |
| Median | 36.0 | 43.9 | 42.0 | 42.0 | 40.0 |
| Range | 19.0–72.0 | 21.5–101.1 | 24.5–84.0 | 19.0–93.5 | 19.0–101.1 |
| BMI, kg/m² | | | | | |
| Mean ± SD | 17.7 ± 3.0 | 20.0 ± 4.1 | 19.2 ± 3.5 | 18.7 ± 3.8 | 18.8 ± 3.6 |
| Median | 17.4 | 19.1 | 18.6 | 18.1 | 18.1 |
| Range | 10.1–29.8 | 11.6–35.0 | 14.0–30.1 | 12.6–32.0 | 10.1–35.0 |

Abbreviations: 9vHPV, 9-valent human papillomavirus; BMI, body mass index; SD, standard deviation.

71.5–98.7) (Table 3); reduced risk of cervical cytological abnormalities related to HPV-52 or HPV-58 was also demonstrated. The incidence of HPV-31/33/45/52/58–related cervical biopsies was reduced by 100% (95% CI, 73.4–100).

Supportive analyses in the HNTS population showed reduction of HPV-31/33/45/52/58–related infection and disease endpoints (Supplementary Table 4). In the mITT analyses, efficacy was demonstrated in susceptible participants (not infected at baseline for the HPV type being analyzed); in the participants infected at baseline, incidence of infection or disease was similar between the 2 vaccine groups (Supplementary Table 4). In the qHPV vaccine group, incidence of persistent infection increased over time in the PPE, HNTS, and mITT populations. In the 9vHPV vaccine group, the incidence of persistent infection was low in the PPE and HNTS populations and started to plateau after month 24 in the mITT population (Supplementary Figure 1).

Immunogenicity

Among Asian young women from Study 001, GMTs for anti-HPV-6/11/16/18 at month 7 were generally similar between the qHPV vaccine and 9vHPV vaccine groups within each country (Table 4). Overall, ≥97.9% of participants underwent seroconversion within 1 month after the last 9vHPV vaccination (ie, study month 7) to each of the vaccine types (Table 4). Among young Indian women in Study 002 (Supplementary Table 5), GMTs were similar or higher across HPV types compared with young women in the overall Study 001 population, indicating that HPV antibody responses in young Indian women were sufficient to induce high-level protective efficacy; seroconversion rates at month 7 were 100% for all 9 HPV types (Supplementary Table 5). However, the number of young Indian women enrolled in Study 002 was limited (N = 25).

Among Asian girls and boys from Study 002, GMTs for each of the 9 HPV types were higher than in the overall population of young women (N = 6792) from Study 001 (Table 5) and the subgroups of young women from each corresponding country in Study 001 (Table 4). Overall, ≥98.8% of Asian girls and boys underwent seroconversion at month 7 to each of the 9 HPV types (Table 5). Results were similar when considering GMTs and seroconversion rates for girls only (Supplementary Table 6).

Safety

In Study 001 (Table 6), injection-site AEs were more common within each country in young women in the 9vHPV vaccine group (85.2%, 81.9%, 87.5%, and 86.6% of participants from Hong Kong/Taiwan, Japan, South Korea, and Thailand, respectively) than in the qHPV vaccine group (77.7%, 79.5%, 82.0%, and 83.1%, respectively). The most common injection-site AEs were pain, swelling, and erythema; most were mild to moderate in intensity. The proportions of participants from Hong Kong/Taiwan, Japan, and South Korea who experienced a vaccine-related systemic AE (9vHPV, 11.8%–18.8%; qHPV, 6.3%–13.9%) were lower than in the overall study population (9vHPV, 29.5%; qHPV, 27.3%) [18, 29]. The proportions of participants from Thailand with vaccine-related systemic AEs (9vHPV, 42.2%; qHPV, 40.3%) were significantly higher than in the overall population (Supplementary Table 7). The frequencies of severe vaccine-related systemic AEs were similar between participants from Thailand and the overall study population (Supplementary Table 7). The difference in frequency of vaccine-related systemic AEs was mainly due to a higher frequency of AEs of pyrexia among participants from Thailand (9vHPV, 16.4%; qHPV, 15.2%) versus the overall population (9vHPV, 5.0%; qHPV, 4.3%).

Table 3. Impact of the 9vHPV Vaccine on the Incidence of HPV-31/33/45/52/58–Related Persistent Infection; Cervical, Vulvar, and Vaginal Disease; Cervical Cytological Abnormalities; and Cervical Medical Procedures in Asian Participants (Young Women Aged 16–26 Years) in the PPE Population in Study 001 by Vaccination Group

| Endpoint | 9vHPV Vaccine (N = 856) | | qHPV Vaccine (N = 857) | | Vaccine Efficacy (95% CI)% |
|--|----------------------------|------|---------------------------|-------|-------------------------------|
| | Cases / n | Rate | Cases / n | Rate | |
| HPV-31/33/45/52/58–related persistent infection ≥6 months duration ^a | 3 / 736 | 13.1 | 67 / 739 | 309.1 | 95.8 (87.8–98.9) |
| By country | | | | | |
| Hong Kong/Taiwan ^b | 0 / 291 | 0.0 | 17 / 300 | 202.1 | 100 (78.2–100) |
| Japan | 2 / 112 | 55.1 | 20 / 117 | 575.7 | 90.4 (62.4–98.4) |
| South Korea | 0 / 124 | 0.0 | 11 / 113 | 354.0 | 100 (71.2–100) |
| Thailand | 1 / 209 | 14.5 | 19 / 209 | 284.2 | 94.9 (70.9–99.8) |
| By HPV type | | | | | |
| HPV-31–related | 0 / 712 | 0.0 | 7 / 702 | 32.5 | 100 (41.0–100) |
| HPV-33–related | 0 / 707 | 0.0 | 8 / 716 | 36.4 | 100 (46.5–100) |
| HPV-45–related | 0 / 726 | 0.0 | 1 / 725 | 4.5 | 100 (≤–999 to 100) |
| HPV-52–related | 3 / 652 | 14.8 | 33 / 644 | 169.0 | 91.3 (74.5–97.7) |
| HPV-58–related | 0 / 666 | 0.0 | 25 / 685 | 120.1 | 100 (86.3–100) |
| HPV-31/33/45/52/58–related persistent infection ≥12 months duration ^c | 3 / 736 | 13.1 | 47 / 739 | 214.1 | 93.9 (81.4–98.4) |
| By HPV type | | | | | |
| HPV-31–related | 0 / 712 | 0.0 | 5 / 702 | 23.2 | 100 (2.5–100) |
| HPV-33–related | 0 / 707 | 0.0 | 8 / 716 | 36.4 | 100 (46.5–100) |
| HPV-45–related | 0 / 726 | 0.0 | 1 / 725 | 4.5 | 100 (≤–999 to 100) |
| HPV-52–related | 3 / 652 | 14.8 | 24 / 644 | 122.0 | 87.9 (61.1–96.9) |
| HPV-58–related | 0 / 666 | 0.0 | 11 / 685 | 52.3 | 100 (66.5–100) |
| HPV-31/33/45/52/58–related cervical, vulvar, and vaginal disease (any grade) | 0 / 751 | 0.0 | 7 / 745 | 29.4 | 100 (40.0–100) |
| By HPV type | | | | | |
| HPV-31–related | 0 / 726 | 0.0 | 0 / 710 | 0.0 | NA |
| HPV-33–related | 0 / 720 | 0.0 | 2 / 722 | 8.7 | 100 (–248.2 to 100) |
| HPV-45–related | 0 / 740 | 0.0 | 0 / 733 | 0.0 | NA |
| HPV-52–related | 0 / 662 | 0.0 | 4 / 650 | 19.2 | 100 (–10.1 to 100) |
| HPV-58–related | 0 / 676 | 0.0 | 1 / 689 | 4.5 | 100 (≤–999 to 100) |
| By lesion type | | | | | |
| HPV-31/33/45/52/58–related cervical disease (any grade) | 0 / 738 | 0.0 | 7 / 737 | 30.5 | 100 (39.7–100) |
| CIN 1 | 0 / 738 | 0.0 | 6 / 737 | 26.2 | 100 (33.0–100) |
| CIN 2/3, AIS, and cervical cancer | 0 / 738 | 0.0 | 1 / 737 | 4.3 | 100 (≤–999 to 100) |
| HPV-31/33/45/52/58–related vulvar and vaginal disease (any grade) | 0 / 751 | 0.0 | 0 / 745 | 0.0 | NA |
| HPV-31/33/45/52/58–related ASC-US HR-HPV positive or worse | 2 / 731 | 8.8 | 25 / 729 | 111.5 | 92.1 (71.5–98.7) |
| By HPV type | | | | | |
| HPV-31–related | 0 / 708 | 0.0 | 3 / 694 | 13.9 | 100 (–68.3 to 100) |
| HPV-33–related | 0 / 703 | 0.0 | 3 / 708 | 13.6 | 100 (–72.8 to 100) |
| HPV-45–related | 0 / 722 | 0.0 | 2 / 717 | 9.0 | 100 (–245.6 to 100) |
| HPV-52–related | 1 / 649 | 4.9 | 15 / 638 | 75.6 | 93.5 (58.4–99.7) |
| HPV-58–related | 1 / 662 | 4.8 | 9 / 678 | 42.8 | 88.7 (18.8–99.5) |
| By lesion type | | | | | |
| ASC-US HR-HPV positive | 1 / 731 | 4.4 | 12 / 729 | 53.0 | 91.7 (51.5–99.6) |
| Low-grade squamous intraepithelial lesion | 1 / 731 | 4.4 | 19 / 729 | 84.5 | 94.8 (70.3–99.7) |
| High-grade squamous intraepithelial lesion or worse ^d | 0 / 731 | 0.0 | 2 / 729 | 8.8 | 100 (–247.0 to 100) |
| HPV-31/33/45/52/58–related cervical biopsy | 0 / 750 | 0.0 | 15 / 745 | 63.8 | 100 (73.4–100) |
| By HPV type | | | | | |
| HPV-31–related | 0 / 725 | 0.0 | 0 / 710 | 0.0 | NA |
| HPV-33–related | 0 / 719 | 0.0 | 2 / 722 | 8.7 | 100 (–248.5 to 100) |
| HPV-45–related | 0 / 739 | 0.0 | 0 / 733 | 0.0 | NA |
| HPV-52–related | 0 / 661 | 0.0 | 11 / 650 | 53.3 | 100 (67.3–100) |
| HPV-58–related | 0 / 675 | 0.0 | 3 / 689 | 13.7 | 100 (–74.7 to 100) |
| HPV-31/33/45/52/58–related cervical definitive therapy | 0 / 750 | 0.0 | 0 / 745 | 0.0 | NA |
| HPV-31/33/45/52/58–related external genital procedures | 0 / 751 | 0.0 | 0 / 745 | 0.0 | NA |

The PPE population consisted of participants who received all 3 doses of vaccine within 1 year, were seronegative at day 1, and had negative results on PCR assays for all HPV types tested from day 1 through month 7 to the vaccine HPV type being analyzed, and had no protocol violations that could interfere with the evaluation of vaccine efficacy as judged by the study director.

Table 3. Continued

Participants are counted once in each applicable endpoint category. A participant may appear in more than 1 category. Rate is the estimate of number of cases per 10000 person-years.

Abbreviations: 9vHPV, 9-valent human papillomavirus; AIS, adenocarcinoma in situ; ASC-US, atypical squamous cells of undetermined significance; CI, confidence interval; CIN, cervical intraepithelial neoplasia; HPV, human papillomavirus; HR, high risk; NA, not available (ie, not calculable); PCR, polymerase chain reaction; PPE, per-protocol efficacy; qHPV, quadrivalent human papillomavirus.

N = number of participants randomized to the respective vaccination group who received at least 1 injection. n = number of participants who have at least 1 follow-up visit after month 7.

^aA case of persistent infection occurred if a participant, after completion of the month 7 visit, is positive for the same HPV type by the HPV-31/33/45/52/58 PCR assay to at least 1 common gene in 2 or more consecutive cervicovaginal/external genital swab, biopsy, or definitive therapy samples obtained at 2 or more consecutive visits at least 6 months (\pm 1 month visit windows) apart.

^bResults for participants from Taiwan only are 0/216 and 13/228 cases in the 9vHPV and qHPV vaccine groups, respectively, representing an efficacy of 100% (95% CI, 71.7–100).

^cA case of persistent infection occurred if a participant, after completion of the month 7 visit, is positive for the same HPV type by the HPV-31/33/45/52/58 PCR assay to at least 1 common gene in 2 or more consecutive cervicovaginal/external genital swab, biopsy, or definitive therapy samples obtained for over a period of at least 12 months (\pm 1 month visit windows) apart.

^dIncludes high-grade squamous intraepithelial lesion; atypical squamous cells, cannot exclude high-grade squamous intraepithelial lesion; atypical glandular cells, adenocarcinoma, and squamous cell carcinoma.

(Supplementary Table 7); AEs of pyrexia experienced by participants from Thailand were mostly mild to moderate (Supplementary Table 7), of short duration (median, 1 day) (Supplementary Table 8), and fevers were mostly low grade (Table 6). AEs of pyrexia and elevated temperatures occurred across all the study sites in Thailand (Supplementary Table 9).

In Study 002 (Table 7), 62.4%, 74.4%, 71.2%, and 85.7% of girls and boys from India, South Korea, Taiwan, and Thailand, respectively, experienced an injection-site AE. The most common injection-site AEs were pain, swelling, and erythema; most were mild to moderate in intensity. Similar results were observed when considering only girls (Supplementary Table 10). The 9vHPV vaccine was generally well tolerated in young Indian women (Supplementary Table 11). Vaccine-related systemic AEs were more common among girls and boys from Thailand (26.4%) than those from India, South Korea, and Taiwan (4.3%–10.8%) (Table 7) but comparable with girls and boys from the overall study population (21.1%) (Supplementary Table 12). The most common vaccine-related AEs were headache and pyrexia. The frequency of vaccine-related pyrexia was higher in girls and boys from Thailand (13.6%) compared with the overall study population (7.2%) (Supplementary Table 12); vaccine-related systemic AEs were mild to moderate (Supplementary Table 12) and generally of short duration (median, 1 day) in girls and boys from Thailand (Supplementary Table 13). AEs of pyrexia occurred at both study sites in Thailand (Supplementary Table 14).

In both studies, discontinuations due to AEs were rare (9vHPV, n = 2; qHPV, n = 1; see Supplementary Material for details). No serious vaccine-related AEs were reported among Asian participants for the entire study duration. Three Asian participants (all from Study 001) died during the entire course of the study (see the Supplementary Material for more information); none of the deaths were considered vaccine related.

DISCUSSION

The 9vHPV vaccine markedly reduced the risk of HPV-31/33/45/52/58-related persistent infection, cervical cytological abnormalities, disease, and medical procedures in Asian participants. Consistent with the HPV type distribution observed in epidemiological studies in Asia, HPV-52

and HPV-58 were most commonly associated with persistent infections and cervical cytological abnormalities in participants who received the qHPV vaccine [1, 6, 31]. In the mITT analyses, nearly all cases of persistent infection and cervical cytological abnormalities and disease occurred among participants who were infected with HPV before vaccination, which highlights the value of implementing vaccination before exposure to HPV. The 9vHPV vaccine induced robust anti-HPV-6/11/16/18/31/33/45/52/58 antibody responses, with seroconversion rates \geq 97.9%. GMTs to HPV-6/11/16/18 were generally comparable between 9vHPV and qHPV vaccine recipients, although GMTs for HPV-11 tended to be lower with 9vHPV vaccine. This is similar to observations in the overall study population and may not be clinically meaningful, as robust HPV-11 antibody responses were observed with both vaccines and no cases of HPV-11-related persistent infection or disease were observed in per-protocol analyses [18, 19]. GMTs to all 9 included HPV types were higher in girls and boys compared with young women from the same country, similar to previous reports in other regions (including Europe, Latin America, and North America [32], and Japan [25]). The 9vHPV vaccine was generally well tolerated in all subgroups. Injection-site AEs were generally more frequent with the 9vHPV vaccine than the qHPV vaccine, as previously observed in the overall study population [18]. Rates of vaccine-related systemic AEs were generally lower among Asian participants than in the overall population. Participants from Thailand, however, were more likely to experience vaccine-related AEs, due to a higher frequency of low-grade, short-lived fevers. The reasons for these differences are unknown; however, their impact is limited, and they do not appear to represent a clinical concern. Overall, the efficacy, immunogenicity, and safety profile of the 9vHPV vaccine was generally consistent with published results from the overall study populations [18–20].

Real-world experience in the 10 years following introduction of qHPV and bivalent HPV vaccines has demonstrated that vaccination prevents HPV infections and HPV-related disease and supports the favorable safety profile of HPV vaccines [9, 33, 34]. A recent meta-analysis found that HPV

Table 4. Summary of Anti-HPV cLIA GMTs and Seropositivity at Month 7 in Asian Participants (Young Women Aged 16–26 Years) in the PPI Population in Study 001 by Vaccination Group and Country

| Assay (cLIA) | 9vHPV Vaccine | | qHPV Vaccine | | 9vHPV Vaccine | | qHPV Vaccine | |
|-----------------------------|---------------|------------------------|--------------|------------------------|---------------|--------------------------|--------------|--------------------------|
| | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % |
| Hong Kong and Taiwan | N = 313 | | N = 312 | | N = 313 | | N = 312 | |
| Anti-HPV-6 | 229 | 734.6 (660.0–817.7) | 237 | 718.0 (646.2–797.7) | 227 | 99.1 (96.9–99.9) | 235 | 99.2 (97.0–99.9) |
| Anti-HPV-11 | 230 | 496.5 (443.1–556.2) | 237 | 616.5 (551.2–689.5) | 229 | 99.6 (97.6–100) | 236 | 99.6 (97.7–100) |
| Anti-HPV-16 | 237 | 2654.6 (2394.3–2943.1) | 245 | 2579.9 (2330.9–2855.5) | 237 | 100 (98.5–100) | 244 | 99.6 (97.7–100) |
| Anti-HPV-18 | 245 | 790.6 (698.5–894.7) | 259 | 669.4 (593.4–755.0) | 244 | 99.6 (97.7–100) | 258 | 99.6 (97.9–100) |
| Anti-HPV-31 | 247 | 607.4 (524.1–703.9) | 260 | 8.5 (7.3–9.8) | 245 | 99.2 (97.1–99.9) | 126 | 48.5 (42.2–54.7) |
| Anti-HPV-33 | 245 | 383.5 (339.2–433.6) | 260 | <4 (<4 to <4) | 242 | 98.8 (96.5–99.7) | 29 | 11.2 (7.6–15.6) |
| Anti-HPV-45 | 250 | 234.6 (206.7–266.2) | 264 | <3 (<3 to <3) | 246 | 98.4 (96.0–99.6) | 24 | 9.1 (5.9–13.2) |
| Anti-HPV-52 | 231 | 313.4 (279.0–352.1) | 238 | <3 (<3 to <3) | 229 | 99.1 (96.9–99.9) | 10 | 4.2 (2.0–7.6) |
| Anti-HPV-58 | 236 | 487.2 (429.4–552.8) | 257 | <4 (<4 to <4) | 234 | 99.2 (97.0, 99.9) | 39 | 15.2 (11.0–20.2) |
| Japan | N = 127 | | N = 127 | | N = 127 | | N = 127 | |
| Anti-HPV-6 | 93 | 839.7 (696.3–1012.7) | 100 | 621.2 (518.5–744.1) | 93 | 100 (96.1–100) | 98 | 98.0 (93.0–99.8) |
| Anti-HPV-11 | 93 | 611.1 (516.4–723.3) | 100 | 607.3 (516.2–714.5) | 93 | 100 (96.1–100) | 100 | 100 (96.4–100) |
| Anti-HPV-16 | 96 | 2672.7 (2269.6–3147.3) | 103 | 2223.4 (1898.7–2603.5) | 96 | 100 (96.2–100) | 103 | 100 (96.5–100) |
| Anti-HPV-18 | 99 | 688.7 (563.3–842.0) | 109 | 471.9 (389.6–571.5) | 99 | 100 (96.3–100) | 106 | 97.2 (92.2–99.4) |
| Anti-HPV-31 | 104 | 672.3 (552.4–818.2) | 109 | 5.9 (4.9–7.2) | 104 | 100 (96.5–100) | 37 | 33.9 (25.1–43.6) |
| Anti-HPV-33 | 109 | 398.3 (344.8–460.2) | 111 | <4 (<4 to <4) | 109 | 100 (96.7–100) | 11 | 9.9 (5.1–17.0) |
| Anti-HPV-45 | 111 | 258.5 (224.8–297.3) | 111 | <3 (<3 to <3) | 111 | 100 (96.7–100) | 6 | 5.4 (2.0–11.4) |
| Anti-HPV-52 | 98 | 306.3 (267.5–350.7) | 96 | <3 (<3 to <3) | 98 | 100 (96.3–100) | 4 | 4.2 (1.1–10.3) |
| Anti-HPV-58 | 95 | 459.6 (399.9–528.3) | 99 | <4 (<4 to <4) | 95 | 100 (96.2–100) | 12 | 12.1 (6.4–20.2) |
| South Korea | N = 152 | | N = 151 | | N = 152 | | N = 151 | |
| Anti-HPV-6 | 94 | 820.1 (684.3–982.8) | 88 | 801.9 (665.1–966.9) | 92 | 97.9 (92.5–99.7) | 88 | 100 (95.9–100) |
| Anti-HPV-11 | 94 | 613.0 (518.3–724.9) | 88 | 658.2 (553.5–782.8) | 94 | 100 (96.2–100) | 88 | 100 (95.9–100) |
| Anti-HPV-16 | 103 | 2641.6 (2323.2–3003.7) | 99 | 2691.6 (2361.0–3068.3) | 103 | 100 (96.5–100) | 99 | 100 (96.3–100) |
| Anti-HPV-18 | 109 | 634.9 (531.0–759.1) | 104 | 627.0 (522.2–752.8) | 108 | 99.1 (95.0–100) | 104 | 100 (96.5–100) |
| Anti-HPV-31 | 113 | 553.9 (455.4–673.8) | 99 | 7.6 (6.2–9.4) | 112 | 99.1 (95.2–100) | 45 | 45.5 (35.4–55.8) |
| Anti-HPV-33 | 112 | 337.2 (297.1–382.6) | 102 | <4 (<4 to <4) | 112 | 100 (96.8–100) | 5 | 4.9 (1.6–11.1) |
| Anti-HPV-45 | 114 | 213.8 (183.0–249.6) | 102 | <3 (<3 to <3) | 113 | 99.1 (95.2–100) | 5 | 4.9 (1.6–11.1) |
| Anti-HPV-52 | 100 | 310.5 (262.0–368.0) | 93 | <3 (<3 to <3) | 100 | 100 (96.4–100) | 3 | 3.2 (0.7–9.1) |
| Anti-HPV-58 | 104 | 420.1 (361.3–488.5) | 99 | <4 (<4 to <4) | 104 | 100 (96.5–100) | 15 | 15.2 (8.7–23.8) |
| Thailand | N = 232 | | N = 233 | | N = 232 | | N = 233 | |
| Anti-HPV-6 | 158 | 910.5 (806.4–1028.0) | 168 | 832.5 (740.0–936.5) | 158 | 100 (97.7–100) | 168 | 100 (97.8–100) |
| Anti-HPV-11 | 158 | 668.1 (586.2–761.4) | 168 | 736.7 (649.0–836.3) | 158 | 100 (97.7–100) | 168 | 100 (97.8–100) |
| Anti-HPV-16 | 172 | 3479.4 (3125.1–3873.9) | 182 | 3371.3 (3037.1–3742.3) | 172 | 100 (97.9–100) | 182 | 100 (98.0–100) |
| Anti-HPV-18 | 181 | 1095.5 (960.7–1249.2) | 183 | 890.7 (781.7–1014.9) | 181 | 100 (98.0–100) | 183 | 100 (98.0–100) |
| Anti-HPV-31 | 195 | 809.1 (687.0–952.9) | 189 | 14.2 (12.0–16.8) | 194 | 99.5 (97.2–100) | 117 | 61.9 (54.6–68.9) |
| Anti-HPV-33 | 189 | 473.3 (415.1–539.7) | 196 | <4 (<4 to <4) | 188 | 99.5 (97.1–100) | 28 | 14.3 (9.7–20.0) |
| Anti-HPV-45 | 200 | 357.8 (313.7–408.0) | 201 | <3 (<3 to <3) | 199 | 99.5 (97.2–100) | 24 | 11.9 (7.8–17.2) |
| Anti-HPV-52 | 172 | 387.7 (343.1–438.2) | 174 | <3 (<3 to <3) | 171 | 99.4 (96.8–100) | 8 | 4.6 (2.0–8.9) |
| Anti-HPV-58 | 177 | 545.0 (481.8–616.4) | 184 | <4 (<4 to 4.2) | 177 | 100 (97.9–100) | 39 | 21.2 (15.5–27.8) |
| Total | N = 824 | | N = 823 | | N = 824 | | N = 823 | |
| Anti-HPV-6 | 574 | 810.9 (756.9–868.7) | 593 | 742.7 (694.1–794.8) | 570 | 99.3 (98.2–99.8) | 589 | 99.3 (98.3–99.8) |
| Anti-HPV-11 | 575 | 576.6 (537.7–618.3) | 593 | 653.1 (609.7–699.6) | 574 | 99.8 (99.0–100) | 592 | 99.8 (99.1–100) |
| Anti-HPV-16 | 608 | 2866.4 (2695.9–3047.8) | 629 | 2738.7 (2578.4–2908.9) | 608 | 100 (99.4–100) | 628 | 99.8 (99.1–100) |
| Anti-HPV-18 | 634 | 817.8 (757.7–882.7) | 655 | 676.9 (628.0–729.7) | 632 | 99.7 (98.9–100) | 651 | 99.4 (98.4–99.8) |
| Anti-HPV-31 | 659 | 661.3 (605.5–722.2) | 657 | 9.1 (8.3–10.0) | 655 | 99.4 (98.5–99.8) | 325 | 49.5 (45.6–53.4) |
| Anti-HPV-33 | 655 | 401.2 (374.8–429.5) | 669 | <4 (<4 to <4) | 651 | 99.4 (98.4–99.8) | 73 | 10.9 (8.7–13.5) |
| Anti-HPV-45 | 675 | 265.9 (247.7–285.5) | 678 | <3 (<3 to <3) | 669 | 99.1 (98.1–99.7) | 59 | 8.7 (6.7–11.1) |
| Anti-HPV-52 | 601 | 331.3 (309.8–354.3) | 601 | <3 (<3 to <3) | 598 | 99.5 (98.5–99.9) | 25 | 4.2 (2.7–6.1) |
| Anti-HPV-58 | 612 | 486.3 (453.8–521.1) | 639 | <4 (<4 to <4) | 610 | 99.7 (98.8–100) | 105 | 16.4 (13.6–19.5) |

The PPI population included all participants who had no protocol violation that could interfere with the evaluation of the immune response to vaccine as judged by the study director, received all 3 vaccinations within acceptable day ranges, were seronegative at day 1 and PCR negative day 1 through month 7 for the relevant HPV type(s), and had a month 7 serum sample collected within an acceptable day range.

Seropositive percent represents proportion of participants with anti-HPV serum levels ≥ 30 , 16, 20, 24, 10, 8, 8, 8, and 8 mMU/mL for HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58, respectively.

Abbreviations: 9vHPV, 9-valent human papillomavirus; CI, confidence interval; cLIA, competitive Luminex immunoassay; GMT, geometric mean titer; mMU, milli-Merck units; PPI, per-protocol immunogenicity; qHPV, quadrivalent human papillomavirus.

N = number of participants randomized to the respective vaccination group who received at least 1 injection. n = number of participants contributing to the analysis. m = number of participants who had seroconversion.

Table 5. Summary of Anti-HPV cLIA GMTs and Seropositivity at Month 7 in the PPI Population of Asian Girls and Boys (Aged 9–15 Years) in Study 002, by Country, and the Overall Population of Young Women (Aged 16–26 Years) From Study 001

| | | Study 002 | | | | | | | | | | Study 001 ^a | | | | | | | | | | | | |
|--------------|-----|--------------------------|-----|--------------------------|-----|--------------------------|------------------|--------------------------|-----|--------------------------|------|--------------------------|---|--------------------------|---|--------------------------|-----------------|--------------------------|---|--------------------------|---|-------------------------------------|---|--------------------------|
| | | South Korea (N = 129) | | | | | Taiwan (N = 139) | | | | | Thailand (N = 140) | | | | | Total (N = 608) | | | | | Overall Study Population (N = 6792) | | |
| Assay (cLIA) | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL | n | GMT (95% CI), mMU/mL |
| Anti-HPV-6 | 170 | 1641.2 (1428.0–1886.2) | 120 | 2226.3 (1931.8–2565.6) | 134 | 1308.8 (1129.9–1516.0) | 131 | 1464.8 (1277.5–1679.6) | 555 | 1615.9 (1503.3–1736.8) | 3993 | 893.1 (871.7–915.1) | | | | | | | | | | | | |
| Anti-HPV-11 | 170 | 1180.0 (1024.2–1359.5) | 120 | 1524.0 (1318.2–1761.9) | 134 | 956.6 (836.9–1093.5) | 131 | 1024.5 (888.4–1181.5) | 555 | 1146.6 (1067.2–1231.9) | 3995 | 666.3 (649.6–683.4) | | | | | | | | | | | | |
| Anti-HPV-16 | 175 | 7061.4 (6125.2–8140.8) | 121 | 8265.5 (7182.8–9511.5) | 135 | 5804.4 (5142.0–6552.1) | 135 | 6598.6 (5742.3–7582.6) | 566 | 6857.8 (6399.4–7349.0) | 4032 | 3131.1 (3057.1–3206.9) | | | | | | | | | | | | |
| Anti-HPV-18 | 178 | 2118.5 (1798.9–2494.9) | 120 | 2530.4 (2147.5–2981.7) | 137 | 1688.0 (1469.3–1939.3) | 132 | 2433.7 (2081.7–2845.1) | 567 | 2150.5 (1986.0–2328.8) | 4539 | 804.6 (782.7–827.1) | | | | | | | | | | | | |
| Anti-HPV-31 | 171 | 1666.0 (1453.4–1909.7) | 119 | 2177.4 (1864.5–2542.9) | 136 | 1636.2 (1424.7–1879.1) | 133 | 1748.3 (1518.0–2013.7) | 559 | 1776.3 (1654.0–1907.6) | 4466 | 658.4 (636.7–680.9) | | | | | | | | | | | | |
| Anti-HPV-33 | 175 | 855.0 (736.9–992.0) | 121 | 1053.1 (907.2–1222.4) | 137 | 796.4 (696.6–910.5) | 134 | 891.4 (770.0–1022.6) | 567 | 887.4 (825.7–953.7) | 4702 | 415.9 (405.6–426.4) | | | | | | | | | | | | |
| Anti-HPV-45 | 179 | 742.0 (627.4–877.5) | 121 | 900.1 (746.6–1085.2) | 137 | 715.2 (602.4–849.0) | 134 | 787.2 (675.7–917.1) | 571 | 776.9 (713.6–845.9) | 4792 | 252.8 (246.2–259.6) | | | | | | | | | | | | |
| Anti-HPV-52 | 176 | 827.0 (708.2–965.8) | 121 | 1005.9 (866.0–1168.5) | 136 | 779.0 (668.6–907.7) | 134 | 753.8 (657.1–864.8) | 567 | 831.6 (771.1–896.9) | 4455 | 379.7 (371.6–388.0) | | | | | | | | | | | | |
| Anti-HPV-58 | 173 | 1198.1 (1030.5–1393.0) | 121 | 1572.7 (1362.8–1814.9) | 136 | 1152.8 (1006.3–1320.7) | 133 | 1209.3 (1045.5–1398.9) | 563 | 1261.2 (1172.3–1357.0) | 4486 | 482.5 (469.9–495.3) | | | | | | | | | | | | |
| Assay (cLIA) | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % | m | Seropositive (95% CI), % |
| Anti-HPV-6 | 168 | 98.8 (95.8–99.9) | 120 | 100 (97.0–100) | 133 | 99.3 (95.9–100) | 131 | 100 (97.2–100) | 552 | 99.5 (98.4–99.9) | 3985 | 99.8 (99.6–99.9) | | | | | | | | | | | | |
| Anti-HPV-11 | 169 | 99.4 (96.8–100) | 120 | 100 (97.0–100) | 134 | 100 (97.3–100) | 131 | 100 (97.2–100) | 554 | 99.8 (99.0–100) | 3994 | 100 (99.9–100) | | | | | | | | | | | | |
| Anti-HPV-16 | 174 | 99.4 (96.9–100) | 121 | 100 (97.0–100) | 135 | 100 (97.3–100) | 135 | 100 (97.3–100) | 565 | 99.8 (99.0–100) | 4031 | 100 (99.9–100) | | | | | | | | | | | | |
| Anti-HPV-18 | 177 | 99.4 (96.9–100) | 120 | 100 (97.0–100) | 137 | 100 (97.3–100) | 132 | 100 (97.2–100) | 566 | 99.8 (99.0–100) | 4532 | 99.8 (99.7–99.9) | | | | | | | | | | | | |
| Anti-HPV-31 | 171 | 100 (97.9–100) | 119 | 100 (96.9–100) | 136 | 100 (97.3–100) | 133 | 100 (97.3–100) | 559 | 100 (99.3–100) | 4457 | 99.8 (99.6–99.9) | | | | | | | | | | | | |
| Anti-HPV-33 | 174 | 99.4 (96.9–100) | 121 | 100 (97.0–100) | 137 | 100 (97.3–100) | 134 | 100 (97.3–100) | 566 | 99.8 (99.0–100) | 4689 | 99.7 (99.5–99.9) | | | | | | | | | | | | |
| Anti-HPV-45 | 178 | 99.4 (96.9–100) | 121 | 100 (97.0–100) | 137 | 100 (97.3–100) | 134 | 100 (97.3–100) | 570 | 99.8 (99.0–100) | 4773 | 99.6 (99.4–99.8) | | | | | | | | | | | | |
| Anti-HPV-52 | 175 | 99.4 (96.9–100) | 121 | 100 (97.0–100) | 136 | 100 (97.3–100) | 134 | 100 (97.3–100) | 566 | 99.8 (99.0–100) | 4446 | 99.8 (99.6–99.9) | | | | | | | | | | | | |
| Anti-HPV-58 | 172 | 99.4 (96.8–100) | 121 | 100 (97.0–100) | 136 | 100 (97.3–100) | 133 | 100 (97.3–100) | 562 | 99.8 (99.0–100) | 4476 | 99.8 (99.6–99.9) | | | | | | | | | | | | |

The PPI population included all participants who had no protocol violation that could interfere with the evaluation of the immune response to vaccine as judged by the study director, received all 3 vaccinations within acceptable day ranges, were seronegative at day 1 and (16 to 26-year-old women only) PCR negative day 1 through month 7 for the relevant HPV type(s), and had a month 7 serum sample collected within an acceptable day range.

Seropositive percent represents the proportion of participants with anti-HPV serum levels ≥ 30 , 16, 20, 24, 10, 8, 8, and 8 mMU/mL for HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58, respectively.

Abbreviations: 9vHPV, 9-valent human papillomavirus; CI, confidence interval; cLIA, competitive Luminex immunoassay; GMT, geometric mean titer; mMU, milli-Merck units; PPI, per-protocol immunogenicity.

N = number of participants randomized to the respective vaccination group who received at least 1 injection. n = number of participants contributing to the analysis. m = number of participants who had seroconversion.

^aBased on [19].

Table 6. AEs in Asian Young Women (Aged 16–26 Years) from Study 001 by Vaccination Group and Country

| Event | 9vHPV Vaccine | | | | | qHPV Vaccine | | | | |
|---|--------------------------------|-----------------|-----------------------|--------------------|-----------------|--------------------------------|-----------------|-----------------------|--------------------|-----------------|
| | Hong Kong and Taiwan (N = 345) | Japan (N = 127) | South Korea (N = 152) | Thailand (N = 232) | Total (N = 856) | Hong Kong and Taiwan (N = 346) | Japan (N = 127) | South Korea (N = 150) | Thailand (N = 231) | Total (N = 854) |
| Participants with 1 or more AE ^a | 304 (88.1) | 108 (85.0) | 137 (90.1) | 211 (90.9) | 760 (88.8) | 297 (85.8) | 103 (81.1) | 130 (86.7) | 206 (89.2) | 736 (86.2) |
| Injection-site event ^b | 294 (85.2) | 104 (81.9) | 133 (87.5) | 201 (86.6) | 732 (85.5) | 269 (77.7) | 101 (79.5) | 123 (82.0) | 192 (83.1) | 685 (80.2) |
| Pain ^c | 286 (82.9) | 104 (81.9) | 131 (86.2) | 199 (85.8) | 720 (84.1) | 260 (75.1) | 100 (78.7) | 119 (79.3) | 189 (81.8) | 668 (78.2) |
| Mild | 216 (62.6) | 63 (49.6) | 76 (50.0) | 103 (44.4) | 458 (53.5) | 217 (62.7) | 76 (59.8) | 86 (57.3) | 93 (40.3) | 472 (55.3) |
| Moderate | 68 (19.7) | 41 (32.3) | 42 (27.6) | 86 (37.1) | 237 (27.7) | 37 (10.7) | 23 (18.1) | 27 (18.0) | 91 (39.4) | 178 (20.8) |
| Severe | 2 (0.6) | 0 | 13 (8.6) | 10 (4.3) | 25 (2.9) | 6 (1.7) | 1 (0.8) | 6 (4.0) | 5 (2.2) | 18 (2.1) |
| Swelling | 160 (46.4) | 57 (44.9) | 79 (52.0) | 71 (30.6) | 367 (42.9) | 99 (28.6) | 53 (41.7) | 53 (35.3) | 63 (27.3) | 268 (31.4) |
| Mild: 0 to ≤2.5 cm | 121 (35.1) | 36 (28.3) | 57 (37.5) | 51 (22.0) | 265 (31.0) | 84 (24.3) | 46 (36.2) | 43 (28.7) | 50 (21.6) | 223 (26.1) |
| Moderate: >2.5 cm to ≤5.0 cm | 27 (7.8) | 14 (11.0) | 11 (7.2) | 15 (6.5) | 67 (7.8) | 12 (3.5) | 5 (3.9) | 6 (4.0) | 11 (4.8) | 34 (4.0) |
| Severe: >5.0 cm | 11 (3.2) | 7 (5.5) | 11 (7.2) | 5 (2.2) | 34 (4.0) | 3 (0.9) | 2 (1.6) | 4 (2.7) | 2 (0.9) | 11 (1.3) |
| Unknown | 1 (0.3) | 0 | 0 | 0 | 1 (0.1) | 0 | 0 | 0 | 0 | 0 |
| Erythema | 126 (36.5) | 51 (40.2) | 59 (38.8) | 26 (11.2) | 262 (30.6) | 88 (25.4) | 48 (37.8) | 36 (24.0) | 15 (6.5) | 187 (21.9) |
| Mild: 0 to ≤2.5 cm | 108 (31.3) | 35 (27.6) | 42 (27.6) | 20 (8.6) | 205 (23.9) | 78 (22.5) | 44 (34.6) | 29 (19.3) | 13 (5.6) | 164 (19.2) |
| Moderate: >2.5 cm to ≤5.0 cm | 13 (3.8) | 11 (8.7) | 10 (6.6) | 5 (2.2) | 39 (4.6) | 9 (2.6) | 3 (2.4) | 5 (3.3) | 2 (0.9) | 19 (2.2) |
| Severe: >5.0 cm | 5 (1.4) | 5 (3.9) | 7 (4.6) | 1 (0.4) | 18 (2.1) | 1 (0.3) | 1 (0.8) | 2 (1.3) | 0 | 4 (0.5) |
| Pruritus ^c | 16 (4.6) | 12 (9.4) | 15 (9.9) | 10 (4.3) | 53 (6.2) | 4 (1.2) | 14 (11.0) | 7 (4.7) | 7 (3.0) | 32 (3.7) |
| Mild | 13 (3.8) | 11 (8.7) | 12 (7.9) | 7 (3.0) | 43 (5.0) | 4 (1.2) | 14 (11.0) | 5 (3.3) | 5 (2.2) | 28 (3.3) |
| Moderate | 3 (0.9) | 1 (0.8) | 2 (1.3) | 3 (1.3) | 9 (1.1) | 0 | 0 | 2 (1.3) | 2 (0.9) | 4 (0.5) |
| Severe | 0 | 0 | 1 (0.7) | 0 | 1 (0.1) | 0 | 0 | 0 | 0 | 0 |
| Systemic event ^d | 136 (39.4) | 41 (32.3) | 70 (46.1) | 128 (55.2) | 375 (43.8) | 141 (40.8) | 38 (29.9) | 72 (48.0) | 139 (60.2) | 390 (45.7) |
| Any vaccine-related systemic event | 65 (18.8) | 15 (11.8) | 19 (12.5) | 98 (42.2) | 197 (23.0) | 48 (13.9) | 8 (6.3) | 19 (12.7) | 93 (40.3) | 168 (19.7) |
| Headache | 10 (2.9) | 5 (3.9) | 4 (2.6) | 40 (17.2) | 59 (6.9) | 9 (2.6) | 4 (3.1) | 3 (2.0) | 41 (17.7) | 57 (6.7) |
| Pyrexia | 9 (2.6) | 4 (3.1) | 2 (1.3) | 38 (16.4) | 53 (6.2) | 6 (1.7) | 0 | 1 (0.7) | 35 (15.2) | 42 (4.9) |
| Serious event | 2 (0.6) | 1 (0.8) | 1 (0.7) | 2 (0.9) | 6 (0.7) | 2 (0.6) | 2 (1.6) | 1 (0.7) | 2 (0.9) | 7 (0.8) |
| Vaccine-related event | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Death | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Discontinuation due to AE ^e | 0 | 0 | 0 | 1 (0.4) | 1 (0.1) | 0 | 0 | 1 (0.7) | 0 | 1 (0.1) |
| Vaccine-related event | 0 | 0 | 0 | 1 (0.4) | 1 (0.1) | 0 | 0 | 1 (0.7) | 0 | 1 (0.1) |
| Serious event | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Serious vaccine-related event | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Participants with temperature data (N) | 344 | 127 | 148 | 231 | 850 | 346 | 126 | 148 | 231 | 851 |
| Maximum temperatures (oral) ^f | | | | | | | | | | |
| ≥37.8°C | 13 (3.8) | 4 (3.1) | 2 (1.4) | 38 (16.5) | 57 (6.7) | 12 (3.5) | 1 (0.8) | 3 (2.0) | 38 (16.5) | 54 (6.3) |
| ≥38.9°C | 1 (0.3) | 0 | 0 | 8 (3.5) | 9 (1.1) | 0 | 0 | 0 | 4 (1.7) | 4 (0.5) |

Every participant is counted a single time for each applicable specific AE.

Abbreviations: 9vHPV, 9-valent human papillomavirus; AE, adverse event; qHPV, quadrivalent human papillomavirus.

N = number of participants who underwent randomization, received at least 1 dose of vaccine, and had at least 1 follow-up visit related to the AE. n = number of participants contributing to the analysis.

^aAEs that were reported within 1 to 15 days after any vaccination.

^bInjection-site events were AEs that were reported within 1 to 5 days after any vaccination. A specific AE appears on this table only if its incidence in 1 or more of the columns is ≥5% incidence after rounding.

^cIntensities of pain and itching were defined as: mild if there was an awareness of the sign or symptom but it did not interfere with usual activities; as moderate if there was enough discomfort to cause interference with usual activity; and as severe if the pain or discomfort was incapacitating, rendering the participant unable to work or carry out usual activities.

^dSystemic events were AEs that were reported within 1 to 15 days after any vaccination. A specific AE appears on this table only if its incidence in 1 or more of the columns is ≥5% incidence after rounding.

^eDiscontinuation due to AE was reported within 1 to 15 days after any vaccination.

^fTemperatures were recorded within 1 to 5 days after any vaccination.

Table 7. AEs in Asian Girls and Boys (Aged 9–15 Years) in Study 002 by Country

| Event | 9vHPV Vaccine | | | | |
|---|-----------------|-----------------------|------------------|--------------------|-----------------|
| | India (N = 194) | South Korea (N = 129) | Taiwan (N = 139) | Thailand (N = 140) | Total (N = 602) |
| | n (%) | n (%) | n (%) | n (%) | n (%) |
| Participants with 1 or more AE ^a | 127 (65.5) | 99 (76.7) | 104 (74.8) | 126 (90.0) | 456 (75.7) |
| Injection-site event ^b | 121 (62.4) | 96 (74.4) | 99 (71.2) | 120 (85.7) | 436 (72.4) |
| Pain ^c | 117 (60.3) | 91 (70.5) | 96 (69.1) | 118 (84.3) | 422 (70.1) |
| Mild | 62 (32.0) | 72 (55.8) | 77 (55.4) | 74 (52.9) | 285 (47.3) |
| Moderate | 46 (23.7) | 17 (13.2) | 17 (12.2) | 39 (27.9) | 119 (19.8) |
| Severe | 9 (4.6) | 2 (1.6) | 2 (1.4) | 5 (3.6) | 18 (3.0) |
| Swelling | 46 (23.7) | 26 (20.2) | 40 (28.8) | 35 (25.0) | 147 (24.4) |
| Mild: 0 to ≤2.5 cm | 35 (18.0) | 12 (9.3) | 29 (20.9) | 23 (16.4) | 99 (16.4) |
| Moderate: >2.5 cm to ≤5.0 cm | 8 (4.1) | 8 (6.2) | 8 (5.8) | 8 (5.7) | 32 (5.3) |
| Severe: >5.0 cm | 3 (1.5) | 6 (4.7) | 3 (2.2) | 4 (2.9) | 16 (2.7) |
| Erythema | 36 (18.6) | 34 (26.4) | 39 (28.1) | 13 (9.3) | 122 (20.3) |
| Mild: 0 to ≤2.5 cm | 31 (16.0) | 27 (20.9) | 33 (23.7) | 9 (6.4) | 100 (16.6) |
| Moderate: >2.5 cm to ≤5.0 cm | 4 (2.1) | 3 (2.3) | 4 (2.9) | 3 (2.1) | 14 (2.3) |
| Severe: >5.0 cm | 1 (0.5) | 4 (3.1) | 2 (1.4) | 1 (0.7) | 8 (1.3) |
| Pruritus ^c | 1 (0.5) | 11 (8.5) | 6 (4.3) | 1 (0.7) | 19 (3.2) |
| Mild | 1 (0.5) | 11 (8.5) | 3 (2.2) | 1 (0.7) | 16 (2.7) |
| Moderate | 0 | 0 | 3 (2.2) | 0 | 3 (0.5) |
| Severe | 0 | 0 | 0 | 0 | 0 |
| Edema ^d | 0 | 19 (14.7) | 0 | 0 | 19 (3.2) |
| Mild | 0 | 12 (9.3) | 0 | 0 | 12 (2.0) |
| Moderate | 0 | 4 (3.1) | 0 | 0 | 4 (0.7) |
| Severe | 0 | 3 (2.3) | 0 | 0 | 3 (0.5) |
| Systemic event ^e | 34 (17.5) | 35 (27.1) | 25 (18.0) | 69 (49.3) | 163 (27.1) |
| Any vaccine-related systemic event | 21 (10.8) | 6 (4.7) | 6 (4.3) | 37 (26.4) | 70 (11.6) |
| Headache | 1 (0.5) | 2 (1.6) | 2 (1.4) | 15 (10.7) | 20 (3.3) |
| Pyrexia | 19 (9.8) | 0 | 3 (2.2) | 19 (13.6) | 41 (6.8) |
| Serious event | 0 | 2 (1.6) | 0 | 0 | 2 (0.3) |
| Vaccine-related event | 0 | 0 | 0 | 0 | 0 |
| Death | 0 | 0 | 0 | 0 | 0 |
| Discontinuation due to AE ^f | 0 | 0 | 0 | 0 | 0 |
| Participants with temperature data (N) | 194 | 129 | 139 | 140 | 602 |
| Maximum temperatures (oral) ^g | | | | | |
| ≥37.8°C | 20 (10.3) | 2 (1.6) | 4 (2.9) | 24 (17.1) | 50 (8.3) |
| ≥38.9°C | 0 | 0 | 1 (0.7) | 5 (3.6) | 6 (1.0) |

Every participant is counted a single time for each applicable specific AE.

Abbreviations: 9vHPV, 9-valent human papillomavirus; AE, adverse event.

N = number of participants who underwent randomization, received at least 1 dose of vaccine, and had at least 1 follow-up visit related to the AE. n = number of participants contributing to the analysis.

^aAEs that were reported within 1 to 15 days after any vaccination.

^bInjection-site events were AEs that were reported within 1 to 5 days after any vaccination. A specific AE appears on this table only if its incidence in 1 or more of the columns is ≥5% incidence after rounding.

^cIntensities of pain and itching were defined as: mild if there was an awareness of the sign or symptom but it did not interfere with usual activities; as moderate if there was enough discomfort to cause interference with usual activity; and as severe if the pain or discomfort was incapacitating, rendering the participant unable to work or carry out usual activities.

^dInjection-site edema was reported in 19 girls and boys from a single study site in South Korea in Study 002. Edema was not reported as an AE at other study sites in South Korea or other countries; moreover, this particular study site did not report any event of injection-site swelling, even though this would have been anticipated because injection-site swelling is one of the most common AEs following vaccination. Reports of injection-site AE of edema at this study site may be due to language-related misinterpretation (ie, even though there are 2 distinct Korean words for swelling and edema, respectively, there is also another word for both swelling and edema).

^eSystemic events were AEs that were reported within 1 to 15 days after any vaccination. A specific AE appears on this table only if its incidence in 1 or more of the columns is ≥5% incidence after rounding.

^fDiscontinuation due to AE was reported within 1 to 15 days after any vaccination.

^gTemperatures were recorded within 1 to 5 days after any vaccination.

vaccines were highly immunogenic and well tolerated in Asian populations, consistent with international HPV vaccination studies [35].

Vaccination programs may represent the most effective prevention strategies in countries without organized cervical cancer screening programs [3, 36]. This is relevant to Asia, which

contributes over half of the global cervical cancer burden, and where comprehensive, organized screening programs have not been widely implemented [4]. The 9vHPV vaccine can provide additional benefits over bivalent and quadrivalent HPV vaccines by expanding coverage to HPV-31/33/45/52/58. The expansion of vaccine coverage to HPV-52/58 is of particular importance in Asia due to the relatively high prevalence of these types (see [Supplementary Material](#)) [1, 12]. The robust reduction of HPV-52- and HPV-58-related persistent infection and cervical cytological abnormalities in 9vHPV vaccine recipients should inform cervical screening algorithms in vaccinated populations in Asia.

HPV vaccination is considered likely to be cost effective in preventing HPV-associated cancers, particularly in developing countries with limited access to other forms of cervical cancer prevention and control [37–39]. Based on studies using Japanese, US and Austrian data, a vaccination program for 9vHPV is expected to improve health outcomes and be cost effective compared with programs implemented with bivalent or qHPV vaccines [40–43].

Although several Asian countries have licensed bivalent or qHPV vaccines, few have implemented national HPV vaccination programs to date [3–5], and only approximately 0.2% of Asian women have received a full course of HPV vaccination, compared with 1.4% worldwide [3]. In Malaysia, a national school-based HPV vaccination program has been in place since 2010 and achieved a high degree of coverage [3–5]. Pilot programs had been implemented in India and Thailand as of 2015 [3]. Issues related to loss of public confidence in vaccination and misperceptions surrounding vaccine safety have emerged in some regions [44–46], despite data from large epidemiological studies and active surveillance programs by national and international organizations, which overwhelmingly support the safety of HPV vaccination [47–49]. As misperceptions regarding vaccine safety and clusters of anxiety-related immunization reactions can have damaging consequences for vaccination programs [46, 50], health care professionals and authorities must be prepared to respond to these challenges in implementing successful vaccination programs [50].

In conclusion, the 9vHPV vaccine is efficacious, immunogenic, and generally well tolerated in clinical studies in Asian girls, boys, and young women, consistent with findings in the global clinical program. These data support widespread vaccination programs in Asia. This information may represent valuable insight for decision-making purposes regarding implementation of vaccination programs/HPV-related disease prevention.

Supplementary Data

Supplementary materials are available at *The Journal of Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. Medical writing support was provided by Erin Bekes, PhD, of Complete Medical Communications and was funded by Merck & Co., Inc., Kenilworth, NJ, USA.

Primary Investigators in Asia, Study V503-001: *Hong-Kong*: Tak-Hong Cheung, Yuen-Sheung Hextan Ngan; *Japan*: Hideki Hanashi, Masaru Sakamoto, Nobuhiro Takeshima, Yuzuru Takeuchi, Wataru Tsunazawa; *South Korea*: Chi-Heum Cho, Byoung Gie Kim, Seung Cheol Kim, Young Tak Kim, Hee-Sug Ryu, Yong Sang Song; *Taiwan*: Chi-An Chen, Hung-Hsueh Chou, Tang-Yuan Chu, Chung-Yuan Lee, Nae-Fang Twu, Yuh-Cheng Yang, Mu-Hsien Yu; *Thailand*: Nipon Khemapech, Sompong Koonlerkit, Sutham Pinjaroen, Punnee Pitisuttithum, Sarikapan Wilailak.

Study V503-002: *India*: Amaresh Adyar, Neerja Bhatla, Sukanta Chatterjee, Hemant Jain, Sanjay Lalwani, Sudha Marwah, Daljit Singh; *Korea*: Chi-Heum Cho, Jin Han Kang, Dongsoo Kim, Kyung-Hyo Kim; *Taiwan*: Li-Min Huang, Tzou-Yien Liu, Horng-Der Tsai; *Thailand*: Punnee Pitisuttithum, Rudiwilai Samakoses.

Financial support. This work was supported by Merck & Co., Inc., Kenilworth, NJ, USA.

Potential conflicts of interest. S. M. G. has received grants from Merck & Co., Inc., Kenilworth, NJ, USA, GlaxoSmithKline, CSL, and Commonwealth Department of Health through her institution, travel, and accommodation expenses paid by Merck & Co., Inc., Kenilworth, NJ, USA, to present at HPV advisory board meetings, and has delivered lectures and received speaking fees from Merck Sharpe & Dohme (MSD) and SPMSD for work performed in her personal time. L.-M. H. has received personal fees as a lecturer for educational symposium from MSD, a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA. N. B. has received research funding through her institution for this study. H. Y. has received lecture and advisory fees from GlaxoSmithKline K.K. and MSD K.K., Tokyo, Japan. M. C. E., E. M., M. R., C. S., A. W., G. P., and A. L. are employees of Merck & Co., Inc., Kenilworth, NJ, USA, and may have received stock/stock options. S. R. H., S. M., and M. S. are employees of MSD K.K., Tokyo, Japan. All other authors report no potential conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Bruni L, Barrinuevo-Rosas L, Albero G, et al. Human papillomavirus and related diseases report: Asia. Summary Report. <http://www.hpvcntr.net/statistics/reports/XSX.pdf>. Accessed 2 August 2017.
2. de Martel C, Plummer M, Vignat J, Franceschi S. Worldwide burden of cancer attributable to HPV by site, country and HPV type. *Int J Cancer* 2017; 141:664–70.
3. Bruni L, Diaz M, Barrionuevo-Rosas L, et al. Global estimates of human papillomavirus vaccination coverage by

- region and income level: a pooled analysis. *Lancet Glob Health* **2016**; 4:e453–63.
4. Garland SM, Bhatla N, Ngan HY. Cervical cancer burden and prevention strategies: Asia Oceania perspective. *Cancer Epidemiol Biomarkers Prev* **2012**; 21:1414–22.
 5. Shin HR, Shin A, Woo H, et al. Prevention of infection-related cancers in the WHO Western Pacific Region. *Jpn J Clin Oncol* **2016**; 46:13–22.
 6. de Sanjose S, Quint WG, Alemany L, et al.; Retrospective International Survey and HPV Time Trends Study Group. Human papillomavirus genotype attribution in invasive cervical cancer: a retrospective cross-sectional worldwide study. *Lancet Oncol* **2010**; 11:1048–56.
 7. Garland SM, Steben M, Sings HL, et al. Natural history of genital warts: analysis of the placebo arm of 2 randomized phase III trials of a quadrivalent human papillomavirus (types 6, 11, 16, and 18) vaccine. *J Infect Dis* **2009**; 199:805–14.
 8. Schiller JT, Castellsagué X, Garland SM. A review of clinical trials of human papillomavirus prophylactic vaccines. *Vaccine* **2012**; 30:F123–38.
 9. Drolet M, Bénard É, Boily MC, et al. Population-level impact and herd effects following human papillomavirus vaccination programmes: a systematic review and meta-analysis. *Lancet Infect Dis* **2015**; 15:565–80.
 10. Malagón T, Drolet M, Boily MC, et al. Cross-protective efficacy of two human papillomavirus vaccines: a systematic review and meta-analysis. *Lancet Infect Dis* **2012**; 12:781–9.
 11. Tanton C, Mesher D, Beddows S, et al. Human papillomavirus (HPV) in young women in Britain: population-based evidence of the effectiveness of the bivalent immunisation programme and burden of quadrivalent and 9-valent vaccine types. *Papillomavirus Res* **2017**; 3:36–41.
 12. Bruni L, Barrinuevo-Rosas L, Albero G, et al. Human papillomavirus and related diseases report: world. Summary Report. <http://www.hpvcentre.net/statistics/reports/XWX.pdf>. Accessed 16 Dec 2016.
 13. Brotherton JML, Tabrizi SN, Phillips S, et al. Looking beyond human papillomavirus (HPV) genotype 16 and 18: defining HPV genotype distribution in cervical cancers in Australia prior to vaccination. *Int J Cancer* **2017**; 141:1576–84.
 14. Alemany L, Saunier M, Alvarado-Cabrero I, et al.; HPV VVAP Study Group. Human papillomavirus DNA prevalence and type distribution in anal carcinomas worldwide. *Int J Cancer* **2015**; 136:98–107.
 15. Joura EA, Ault KA, Bosch FX, et al. Attribution of 12 high-risk human papillomavirus genotypes to infection and cervical disease. *Cancer Epidemiol Biomarkers Prev* **2014**; 23:1997–2008.
 16. Serrano B, de Sanjosé S, Tous S, et al. Human papillomavirus genotype attribution for HPVs 6, 11, 16, 18, 31, 33, 45, 52 and 58 in female anogenital lesions. *Eur J Cancer* **2015**; 51:1732–41.
 17. Serrano B, Alemany L, Tous S, et al. Potential impact of a nine-valent vaccine in human papillomavirus related cervical disease. *Infect Agent Cancer* **2012**; 7:38.
 18. Joura EA, Giuliano AR, Iversen OE, et al.; Broad Spectrum HPV Vaccine Study. A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women. *N Engl J Med* **2015**; 372:711–23.
 19. Huh WK, Joura EA, Giuliano AR, et al. Final efficacy, immunogenicity, and safety analyses of a nine-valent human papillomavirus vaccine in women aged 16–26 years: a randomised, double-blind trial. *Lancet* **2017**; 390:2143–59.
 20. Van Damme P, Olsson SE, Block S, et al. Immunogenicity and safety of a 9-valent HPV vaccine. *Pediatrics* **2015**; 136:e28–39.
 21. Chen YH, Gesser R, Luxembourg A. A seamless phase IIB/III adaptive outcome trial: design rationale and implementation challenges. *Clin Trials* **2015**; 12:84–90.
 22. Luxembourg A, Bautista O, Moeller E, Ritter M, Chen J. Design of a large outcome trial for a multivalent human papillomavirus L1 virus-like particle vaccine. *Contemp Clin Trials* **2015**; 42:18–25.
 23. Luxembourg A, Brown D, Bouchard C, et al. Phase II studies to select the formulation of a multivalent HPV L1 virus-like particle (VLP) vaccine. *Hum Vaccin Immunother* **2015**; 11:1313–22.
 24. Luxembourg A, Moreira ED Jr, Samakoses R, et al. Phase III, randomized controlled trial in girls 9–15 years old to evaluate lot consistency of a novel nine-valent human papillomavirus L1 virus-like particle vaccine. *Hum Vaccin Immunother* **2015**; 11:1306–12.
 25. Iwata S, Murata S, Rong Han S, Wakana A, Sawata M, Tanaka Y. Safety and immunogenicity of a 9-valent human papillomavirus vaccine administered to 9- to 15-year-old Japanese girls. *Jpn J Infect Dis* **2017**; 70:368–73.
 26. Else EA, Swoyer R, Zhang Y, et al. Comparison of real-time multiplex human papillomavirus (HPV) PCR assays with INNO-LiPA HPV genotyping extra assay. *J Clin Microbiol* **2011**; 49:1907–12.
 27. Roberts CC, Swoyer R, Bryan JT, Taddeo FJ. Comparison of real-time multiplex human papillomavirus (HPV) PCR assays with the linear array HPV genotyping PCR assay and influence of DNA extraction method on HPV detection. *J Clin Microbiol* **2011**; 49:1899–906.
 28. Roberts C, Green T, Hess E, et al. Development of a human papillomavirus competitive luminex immunoassay for 9 HPV types. *Hum Vaccin Immunother* **2014**; 10:2168–74.
 29. Moreira ED, Jr, Block SL, Ferris D, et al. Safety profile of the 9-valent HPV vaccine: a combined analysis of 7 phase III clinical trials. *Pediatrics* **2016**; 138:e20154387.

30. Chan ISF, Bohidar NR. Exact power and sample size for vaccine efficacy studies. *Comm Stat Theory Methods* **1998**; 27:1305–22.
31. Chan PK, Ho WC, Yu MY, et al. Distribution of human papillomavirus types in cervical cancers in Hong Kong: current situation and changes over the last decades. *Int J Cancer* **2009**; 125:1671–7.
32. Petersen LK, Restrepo J, Moreira Jr ED, et al. Impact of baseline covariates on the immunogenicity of the 9-valent HPV vaccine – a combined analysis of five phase III clinical trials. *Papillomavirus Res* **2017**; 3:105–15.
33. Garland SM, Kjaer SK, Muñoz N, et al. Impact and effectiveness of the quadrivalent human papillomavirus vaccine: a systematic review of 10 years of real-world experience. *Clin Infect Dis* **2016**; 63:519–27.
34. Vichnin M, Bonanni P, Klein NP, et al. An overview of quadrivalent human papillomavirus vaccine safety: 2006 to 2015. *Pediatr Infect Dis J* **2015**; 34:983–91.
35. Setiawan D, Luttjeboer J, Pouwels KB, Wilschut JC, Postma MJ. Immunogenicity and safety of human papillomavirus (HPV) vaccination in Asian populations from six countries: a meta-analysis. *Jpn J Clin Oncol* **2017**; 47:265–76.
36. Pitisuttithum P, Velicer C, Luxembourg A. 9-Valent HPV vaccine for cancers, pre-cancers and genital warts related to HPV. *Expert Rev Vaccines* **2015**; 14:1405–19.
37. Fesenfeld M, Hutubessy R, Jit M. Cost-effectiveness of human papillomavirus vaccination in low and middle income countries: a systematic review. *Vaccine* **2013**; 31:3786–804.
38. Jit M, Brisson M, Portnoy A, Hutubessy R. Cost-effectiveness of female human papillomavirus vaccination in 179 countries: a PRIME modelling study. *Lancet Glob Health* **2014**; 2:e406–14.
39. Kim SY, Sweet S, Chang J, Goldie SJ. Comparative evaluation of the potential impact of rotavirus versus HPV vaccination in GAVI-eligible countries: a preliminary analysis focused on the relative disease burden. *BMC Infect Dis* **2011**; 11:174.
40. Boiron L, Joura E, Largeron N, Prager B, Uhart M. Estimating the cost-effectiveness profile of a universal vaccination programme with a nine-valent HPV vaccine in Austria. *BMC Infect Dis* **2016**; 16:153.
41. Chesson HW, Markowitz LE, Hariri S, Ekwueme DU, Saraiya M. The impact and cost-effectiveness of nonavalent HPV vaccination in the United States: estimates from a simplified transmission model. *Hum Vaccin Immunother* **2016**; 12:1363–72.
42. Durham DP, Ndeffo-Mbah ML, Skrip LA, Jones FK, Bauch CT, Galvani AP. National- and state-level impact and cost-effectiveness of nonavalent HPV vaccination in the United States. *Proc Natl Acad Sci U S A* **2016**; 113:5107–12.
43. Weiss T, Pillsbury M, Abe M, Sato M, Yamabe K. Projecting the potential public health impact of a 9-valent HPV vaccine in Japan. *Value Health* **2014**; 17:A87–88.
44. Hanley SJ, Yoshioka E, Ito Y, Kishi R. HPV vaccination crisis in Japan. *Lancet* **2015**; 385:2571.
45. Tanaka Y, Ueda Y, Egawa-Takata T, Yagi A, Yoshino K, Kimura T. Outcomes for girls without HPV vaccination in Japan. *Lancet Oncol* **2016**; 17:868–9.
46. World Health Organization (WHO). Global advisory committee on vaccine safety, 2–3 December 2015. *Wkly Epidemiol Rec* **2016**; 91:21–32.
47. Garland SM, Stanley M, Brotherton JM, et al. IPVS policy statement on safety of HPV vaccines. *Papillomavirus Res* **2016**; 2:9–10.
48. World Health Organization (WHO). Human papillomavirus vaccines: WHO position paper. *Wkly Epidemiol Rec* **2017**; 92:241–68.
49. Global Advisory Committee on Vaccine Safety, World Health Organization (WHO). Statement on safety of HPV vaccines. http://www.who.int/vaccine_safety/committee/GACVS_HPVS_statement_17Dec2015.pdf?ua=1. Accessed 20 June 2017.
50. World Health Organization (WHO). Guide to introducing HPV vaccine into national immunization programmes. <http://apps.who.int/iris/bitstream/10665/253123/1/9789241549769-eng.pdf?ua=1>. Accessed 15 March 2017.