

Original Article
Endocrinology,
Nutrition & Metabolism



The Association of Adiponectin and Visceral Fat with Insulin Resistance and β -Cell Dysfunction

Hyun Uk Moon ,¹ Kyoung Hwa Ha ,^{1,2} Seung Jin Han ,^{1,2} Hae Jin Kim ,^{1,2} and Dae Jung Kim ,^{1,2,3}

¹Department of Endocrinology and Metabolism, Ajou University School of Medicine, Suwon, Korea
²Cardiovascular and Metabolic Disease Etiology Research Center, Ajou University School of Medicine, Suwon, Korea
³Institute on Aging, Ajou University Medical Center, Suwon, Korea



Received: Apr 18, 2018

Accepted: Nov 8, 2018

Address for Correspondence:

Dae Jung Kim, MD

Department of Endocrinology and Metabolism, Ajou University School of Medicine, 164 World Cup-ro, Yeongtong-gu, Suwon 16499, Republic of Korea.
E-mail: djkim@ajou.ac.kr

© 2019 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Hyun Uk Moon
<https://orcid.org/0000-0002-8772-2932>

Kyoung Hwa Ha
<https://orcid.org/0000-0002-3408-7568>

Seung Jin Han
<https://orcid.org/0000-0003-4783-6799>

Hae Jin Kim
<https://orcid.org/0000-0002-8958-7164>

Dae Jung Kim
<https://orcid.org/0000-0003-1025-2044>

Funding

This research was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health and Welfare, Republic of Korea (HI13C0715).

ABSTRACT

Background: Obesity is a risk factor for metabolic abnormalities. We investigated the relationship of adiponectin levels and visceral adiposity with insulin resistance and β -cell dysfunction.

Methods: This cross-sectional study enrolled 1,347 participants (501 men and 846 women aged 30–64 years) at the Cardiovascular and Metabolic Diseases Etiology Research Center. Serum adiponectin levels and visceral fat were measured using enzyme-linked immunosorbent assay kits and dual-energy X-ray absorptiometry, respectively. Insulin resistance was evaluated using the homeostatic model assessment of insulin resistance (HOMA-IR) and Matsuda insulin sensitivity index. β -cell dysfunction was evaluated using the homeostatic model assessment of β -cell function (HOMA- β), insulinogenic index, and disposition index.

Results: Regarding insulin resistance, compared with individuals with the highest adiponectin levels and visceral fat mass < 75th percentile, the fully adjusted odds ratios (ORs) for HOMA-IR \geq 2.5 and Matsuda index < 25th percentile were 13.79 (95% confidence interval, 7.65–24.83) and 8.34 (4.66–14.93), respectively, for individuals with the lowest adiponectin levels and visceral fat \geq 75th percentile. Regarding β -cell dysfunction, the corresponding ORs for HOMA- β < 25th percentile, insulinogenic index < 25th percentile, and disposition index < 25th percentile were 1.20 (0.71–2.02), 1.01 (0.61–1.66), and 1.87 (1.15–3.04), respectively.

Conclusion: Low adiponectin levels and high visceral adiposity might affect insulin resistance and β -cell dysfunction.

Keywords: Adiponectin; Insulin Resistance; Insulin Secretion; Visceral Fat

INTRODUCTION

Obesity has become a major health concern worldwide and is a major risk factor for insulin resistance and type 2 diabetes.¹ Moreover, excess adipose tissue, in particular visceral adipose tissue (VAT), is an important risk factor for insulin resistance, which links abdominal VAT to cardiovascular risk.² Abdominal VAT could cause insulin resistance by secreting inflammatory adipokines such as interleukin-6, tumor necrosis factor- α , macrophage chemoattractant protein-1, and resistin.³

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Moon HU, Ha KH, Han SJ, Kim HJ, Kim DJ. Data curation: Ha KH. Formal analysis: Ha KH. Investigation: Moon HU, Ha KH. Methodology: Ha KH, Han SJ, Kim HJ, Kim DJ. Software: Ha KH. Supervision: Han SJ, Kim HJ, Kim DJ. Validation: Ha KH. Visualization: Ha KH, Moon HU. Writing - original draft: Moon HU. Writing - review & editing: Ha KH, Han SJ, Kim HJ, Kim DJ.

Adiponectin, which is the most abundant anti-inflammatory and atheroprotective adipokine secreted by adipose tissues, reduces hepatic gluconeogenesis and increases glucose uptake and whole-body insulin sensitivity.⁴ However, plasma adiponectin levels are low in individuals with visceral obesity.^{5,6} In addition, a reduction in plasma adiponectin levels may play a key role in the development of insulin resistance.³ A recent study suggested that inflammatory cytokine-induced DNA methyltransferase 1 expression alters adiponectin expression by selectively methylating and stimulating a compact chromatin structure in the adiponectin promoter.⁷ In addition, a previous study reported that the anti-diabetic effects of adiponectin affect insulin sensitivity and have a direct effect on insulin secretion.⁸

Many studies have shown that adiponectin and visceral fat could be used to predict metabolic abnormalities.⁹⁻¹² However, in most studies, these two factors played a key role in insulin resistance,¹³⁻¹⁷ and only two studies showed a strong association between the combined effects of low adiponectin levels and a high visceral fat mass and the risk of metabolic syndrome in Asian populations.^{9,10} As far as we know, there has been no research that assessed the correlation between the two metabolic factors and two insulin status (insulin resistance and insulin secretion) at the same time, specifically in Asia. Thus, we evaluated the association of adiponectin levels and visceral fat mass with the risk of insulin resistance and β -cell dysfunction.

METHODS**Study population**

Data from the Cardiovascular and Metabolic Diseases Etiology Research Center (CMERC) study, which was initiated in 2013, were used. The CMERC study recruited individuals aged 30–64 years who had no prior history of malignant tumors or cardiovascular diseases. Details of the CMERC study, and the methods used therein, have been described previously.¹⁸ Among the 1,574 participants enrolled from January 2014 to May 2016 in three districts (Suwon, Yongin, and Hwaseong in Korea), 277 were excluded due to lacking data such as VAT mass or 2 hours post-load insulin level. Ultimately, 1,347 individuals were selected for statistical analysis.

Measurements

All participants were interviewed by trained personnel using standardized questionnaire surveys according to a predefined protocol. The standardized questionnaire was used to ascertain basic demographic data, medical history, and information regarding personal characteristics such as physical activity. The level of physical activity was divided into three categories based on the Korean version of International Physical Activity Questionnaire-Short Form: low, moderate, and high physical activity.

Height (cm) and weight (kg) were measured using an automatic height-weight scale (BSM330; InBody Co., Ltd., Seoul, Korea). Body mass index (BMI) was calculated using the following formula: $\text{weight}/\text{height}^2$ (kg/m^2). Waist circumference (cm) was measured midway between the bottom edge of the last rib and the iliac crest in the mid-axillary plane using a measuring tape (Seca GmbH, Hamburg, Germany). Blood pressure (BP) was measured using an automated BP monitor (HEM-7080IC; Omron Healthcare, Lake Forest, IL, USA). Measurements were taken three times at 2 minutes intervals and the last two of the three measurements were averaged. Whole-body dual-energy X-ray absorptiometry (DXA) scans were acquired using a Lunar iDXA (GE Healthcare, Madison, WI, USA) and analyzed with enCORE software (GE Healthcare). VAT mass (kg) was calculated automatically for the DXA

android region-of-interest with a caudal limit at the top of the iliac crest¹⁹; it was divided into gender-specific VAT masses of < 75th percentile and ≥ 75th percentile.

Blood samples were collected from each participant after at least 8 hours of fasting. Oral glucose tolerance tests (OGTTs) were performed to allow accurate evaluation of glucose metabolism. Each participant drank a 75 g glucose solution, and blood samples were taken at 30, 60, and 120 minutes to measure serum glucose and insulin levels. Biochemical assays were performed at a central laboratory (Seoul Clinical Laboratories, Seoul, Korea). Serum adiponectin levels were measured by enzyme-linked immunosorbent assay kits (R&D Systems, Minneapolis, MN, USA). Glucose levels were obtained using a colorimetric method; total cholesterol, high-density lipoprotein cholesterol, and triglyceride levels were measured enzymatically. High-sensitivity C-reactive protein (hs-CRP) levels were obtained via a turbidimetric immunoassay method using an autoanalyzer (ADVIA 1800 Auto Analyzer; Siemens Medical Solutions, Malvern, PA, USA). Insulin levels were measured using radioimmunoassays (SR 300; STRATEC, Birkenfeld, Germany).

Definitions of insulin resistance and β -cell dysfunction

Insulin resistance was evaluated using homeostatic model assessment of insulin resistance (HOMA-IR) and the Matsuda insulin sensitivity index. Insulin secretion was evaluated using homeostatic model assessment of β -cell function (HOMA- β), insulinogenic index, and disposition index. HOMA-IR was calculated as fasting insulin ($\mu\text{U/mL}$) \times fasting glucose/405 (mg/dL).²⁰ The Matsuda insulin sensitivity index was calculated as $10,000/(\text{fasting glucose [mg/dL]} \times \text{fasting insulin } [\mu\text{U/mL}] \times \text{mean glucose during OGTT [mg/dL]} \times \text{mean insulin during OGTT } [\mu\text{U/mL}])^{0.5}$.²¹ HOMA- β was calculated as $360 \times \text{fasting insulin } (\mu\text{U/mL})/\text{fasting glucose (mg/dL)} - 63$.^{20,22} The insulinogenic index was calculated as the increase in insulin levels from 0 to 30 minutes divided by the increase in glucose levels from 0 to 30 minutes during the OGTT.²³ The disposition index was calculated as the insulinogenic index \times 1/fasting insulin.²⁴ Insulin resistance was defined as a HOMA-IR value ≥ 2.5 ²⁵ or a gender-specific Matsuda index < 25th percentile. β -cell dysfunction was defined as a gender-specific HOMA- β < 25th percentile, insulinogenic index < 25th percentile, or disposition index < 25th percentile.

Statistical analysis

All normally distributed data are presented as means \pm standard deviation (SD). Non-normally distributed data are presented as medians and interquartile ranges. Differences were analyzed using Student's t-tests, or Mann-Whitney U tests for continuous variables, or χ^2 tests for categorical variables. Conditional logistic regression models were used to calculate odds ratios (ORs) and 95% confidence intervals (CIs) for insulin resistance and β -cell dysfunction, with adjustment for the following potential confounders: gender, age, BMI, hs-CRP, triglyceride, and physical activity. Participants were divided into tertiles according to gender-specific adiponectin levels.

To evaluate the association of adiponectin and visceral fat levels with insulin resistance and β -cell dysfunction, participants were divided as follows: group 1, highest tertile of adiponectin and a VAT mass < 75th percentile; group 2, middle tertile of adiponectin and VAT mass < 75th percentile; group 3, lowest tertile of adiponectin and VAT mass < 75th percentile; group 4, highest tertile of adiponectin and VAT mass \geq 75th percentile; group 5, middle tertile of adiponectin and VAT mass \geq 75th percentile; and group 6, lowest tertile of adiponectin and VAT mass \geq 75th percentile. All analyses were performed using SAS software (ver. 9.4; SAS Institute Inc., Cary, NC, USA).

Ethics statement

The study protocol was reviewed and approved by the Institutional Review Board (IRB) of Ajou University (IRB No. AJIRB-BMR-SUR-13-272). All participants provided written informed consent before participation.

RESULTS

The baseline characteristics of the study participants are summarized in **Table 1**. Of the 1,347 participants, 501 (37.2%) were men and 846 (62.8%) were women. The mean \pm SD age of the subjects was 50.7 \pm 8.5 years for men and 50.0 \pm 8.3 years for women. The median (25th–75th percentile) adiponectin level was 3.2 μ g/mL (2.1–4.8 μ g/mL) for men and 5.6 μ g/mL (3.5–8.4 μ g/mL) for women. The median (25th–75th percentile) VAT mass was 1.2 kg (0.7–1.6 kg) for men and 0.6 kg (0.4–0.9 kg) for women. The prevalence of diabetes was 12.8% for men and 7.8% for women.

Table 2 shows the association of adiponectin levels and visceral fat mass with insulin resistance. In an unadjusted model, compared with the highest tertile adiponectin level,

Table 1. Characteristics of the study participants: the CMERC study

Characteristics	Total (n = 1,347)	Men (n = 501)	Women (n = 846)	P value ^a
Age, yr	50.3 \pm 8.4	50.7 \pm 8.5	50.0 \pm 8.3	0.156
BMI, kg/m ²	24.4 \pm 3.1	25.2 \pm 3.0	23.9 \pm 3.0	< 0.001
Waist circumference, cm	82.9 \pm 9.1	88.2 \pm 7.8	79.8 \pm 8.3	< 0.001
VAT mass, kg	0.8 (0.5–1.2)	1.2 (0.7–1.6)	0.6 (0.4–0.9)	< 0.001
Systolic BP, mmHg	119.3 \pm 15.4	124.4 \pm 13.5	116.2 \pm 15.7	< 0.001
Diastolic BP, mmHg	75.5 \pm 10.2	79.6 \pm 9.5	73.0 \pm 9.8	< 0.001
Adiponectin, μ g/mL	4.5 (2.8–7.3)	3.2 (2.1–4.8)	5.6 (3.5–8.4)	< 0.001
Fasting glucose, mg/dL	93.0 (86.0–101.0)	96.0 (89.0–106.0)	91.0 (85.0–99.0)	< 0.001
30 min glucose	152.0 (131.0–175.0)	160.0 (139.0–182.0)	147.0 (126.0–169.0)	< 0.001
2 hr glucose	110.0 (90.0–131.0)	110.0 (87.0–135.0)	109.0 (93.0–129.0)	0.715
Fasting insulin, μ IU/mL	8.2 (6.6–10.5)	8.4 (6.7–10.8)	8.1 (6.5–10.3)	0.178
30 min insulin	43.3 (29.5–63.7)	42.6 (28.5–61.8)	43.6 (30.5–64.9)	0.595
2 hr insulin	31.0 (20.0–47.8)	28.3 (16.6–47.0)	33.1 (22.4–48.2)	0.001
Measures of insulin secretion				
HOMA- β	100.5 (76.5–133.5)	90.5 (70.1–123.4)	106.8 (80.8–139.0)	< 0.001
Insulinogenic index	80.7 (46.2–139.9)	71.2 (41.9–120.6)	86.5 (49.1–154.6)	< 0.001
Disposition index	1.4 (0.8–2.4)	1.3 (0.7–2.1)	1.5 (0.9–2.69)	< 0.001
Measures of insulin sensitivity				
HOMA-IR	1.9 (1.5–2.6)	2.0 (1.6–2.7)	1.8 (1.4–2.5)	0.002
Matsuda index	6.1 (4.4–8.0)	5.9 (4.4–7.8)	6.3 (4.4–8.3)	0.051
Total cholesterol, mg/dL	190.4 \pm 34.3	190.1 \pm 32.8	190.7 \pm 35.1	0.759
HDL cholesterol, mg/dL	53.5 \pm 13.3	47.8 \pm 11.0	56.9 \pm 13.4	< 0.001
Triglycerides, mg/dL	112.0 (79.0–162.0)	137.0 (97.0–189.0)	101.0 (73.0–142.0)	< 0.001
hs-CRP, mg/dL	0.6 (0.3–1.2)	0.7 (0.4–1.4)	0.5 (0.3–1.0)	< 0.001
Diabetes mellitus, No. (%)	130 (9.6)	64 (12.8)	66 (7.8)	0.004
Physical activity, No. (%)				
Low	744 (55.2)	257 (51.3)	487 (57.6)	< 0.001
Medium	460 (34.2)	167 (33.3)	293 (34.6)	
High	143 (10.6)	77 (15.4)	66 (7.8)	

Data are means \pm SD or median (25th–75th percentile) unless otherwise indicated.

CMERC = Cardiovascular and Metabolic Diseases Etiology Research Center, BMI = body mass index, VAT = visceral adipose tissue, BP = blood pressure, HOMA- β = homeostatic model assessment of β -cell function, HOMA-IR = homeostatic model assessment of insulin resistance, HDL = high-density lipoprotein, hs-CRP = high-sensitivity C-reactive protein, SD = standard deviation.

^aP values for comparison between men and women were tested by Student's t-tests, Mann-Whitney U tests, or χ^2 tests.

Table 2. ORs (95% CI) for insulin resistance

Variables	HOMA-IR \geq 2.5				Matsuda index < 25th percentile			
	Unadjusted		Adjusted ^a		Unadjusted		Adjusted ^a	
	OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
Total								
Adiponectin (tertile)								
High	1.00		1.00		1.00		1.00	
Middle	2.60 (1.83–3.69)	< 0.001	1.80 (1.21–2.68)	0.004	2.16 (1.51–3.10)	< 0.001	1.35 (0.90–2.04)	0.152
Low	5.47 (3.90–7.67)	< 0.001	3.34 (2.28–4.91)	< 0.001	4.51 (3.20–6.35)	< 0.001	2.38 (1.61–3.54)	< 0.001
VAT mass \geq 75th percentile (ref: < 75th percentile)	9.36 (7.07–12.37)	< 0.001	3.60 (2.48–5.23)	< 0.001	10.24 (7.69–13.64)	< 0.001	3.44 (2.36–5.04)	< 0.001
Men								
Adiponectin (tertile)								
High	1.00		1.00		1.00		1.00	
Middle	1.59 (0.96–2.63)	0.071	1.11 (0.62–1.97)	0.729	1.31 (0.76–2.24)	0.326	0.72 (0.37–1.38)	0.320
Low	3.09 (1.90–5.03)	< 0.001	2.11 (1.21–3.67)	0.009	2.60 (1.57–4.32)	< 0.001	1.48 (0.80–2.75)	0.215
VAT mass \geq 75th percentile (ref: < 75th percentile)	7.51 (4.81–11.74)	< 0.001	3.77 (2.05–6.94)	< 0.001	8.84 (5.58–14.01)	< 0.001	2.64 (1.38–5.05)	0.003
Women								
Adiponectin (tertile)								
High	1.00		1.00		1.00		1.00	
Middle	4.20 (2.50–7.06)	< 0.001	2.93 (1.64–5.21)	< 0.001	3.23 (1.96–5.34)	< 0.001	2.07 (1.18–3.62)	0.011
Low	9.30 (5.62–15.38)	< 0.001	5.67 (3.21–10.01)	< 0.001	6.94 (4.29–11.24)	< 0.001	3.61 (2.09–6.23)	< 0.001
VAT mass \geq 75th percentile (ref: < 75th percentile)	11.04 (7.69–15.86)	< 0.001	3.47 (2.14–5.61)	< 0.001	11.26 (7.80–16.25)	< 0.001	3.90 (2.41–6.33)	< 0.001

Values are presented as median (interquartile range) or number (%).

OR = odds ratio, CI = confidence interval, HOMA-IR = homeostatic model assessment of insulin resistance, VAT = visceral adipose tissue, BMI = body mass index, hs-CRP = high-sensitivity C-reactive protein.

^aAdjusted for gender, age, BMI, hs-CRP, triglyceride, physical activity, adiponectin level, and VAT mass.

the ORs for HOMA-IR \geq 2.5 were 2.60 (95% CI, 1.83–3.69) and 5.47 (3.90–7.67) for the middle and lowest tertile of adiponectin level, respectively. The ORs (95% CI) for a Matsuda index \leq 25th percentile were 2.16 (1.51–3.10) and 4.51 (3.20–6.35) for the middle and lowest tertile, respectively. The association remained significant after adjustment for potential confounders, except the OR for Matsuda index < 25th percentile for the middle tertile. After adjustment for confounders, a VAT mass \geq 75th percentile was associated with a significantly higher risk of insulin resistance (HOMA-IR \geq 2.5: OR, 3.60; 95% CI, 2.48–5.23; Matsuda index < 25th percentile: OR, 3.44; 95% CI, 2.36–5.04) compared with a VAT mass < 75th percentile. These associations were more obvious in women.

Table 3 shows the association of adiponectin level and visceral fat with β -cell dysfunction. After adjustment for confounders, there was no significant association between adiponectin level or visceral fat mass and a HOMA- β < 25th percentile and insulinogenic index < 25th percentile. However, a VAT mass \geq 75th percentile was associated with a significantly higher risk of a disposition index < 25th percentile (OR, 1.70; 95% CI, 1.19–2.42) compared with a VAT mass < 75th percentile. The associations were consistent in both men and women.

The combination of the lowest adiponectin levels and a VAT mass \geq 75th percentile was associated with a higher insulin resistance (HOMA-IR \geq 2.5: adjusted OR, 13.79; 95% CI, 7.65–24.83; Matsuda index < 25th percentile: adjusted OR, 8.34; 95% CI, 4.66–14.93) compared with the highest adiponectin levels and a VAT mass < 75th percentile. In addition, the lowest adiponectin levels and VAT mass \geq 75th percentile showed a stronger association with a disposition index < 25th percentile (OR, 1.87; 95% CI, 1.15–3.04) than did the highest adiponectin levels and VAT mass < 75th percentile in both men and women (**Fig. 1** and **Supplementary Fig. 1**). However, regarding β -cell dysfunction, the associations were

Table 3. ORs (95% CI) for β -cell dysfunction

Variables	HOMA- β < 25th percentile			Insulinogenic index < 25th percentile			Disposition index < 25th percentile			
	Unadjusted OR (95% CI)	P value	Adjusted ^a OR (95% CI)	Unadjusted OR (95% CI)	P value	Adjusted ^a OR (95% CI)	Unadjusted OR (95% CI)	P value	Adjusted ^a OR (95% CI)	P value
Total										
Adiponectin (tertile)	1.00		1.00	1.00		1.00	1.00		1.00	
High	0.93 (0.68-1.25)	0.616	1.08 (0.78-1.50)	0.82 (0.61-1.11)	0.203	0.80 (0.58-1.10)	1.16 (0.86-1.58)	0.332	0.97 (0.70-1.34)	0.860
Middle	1.02 (0.76-1.38)	0.877	1.23 (0.88-1.71)	0.96 (0.71-1.29)	0.764	0.91 (0.66-1.25)	1.54 (1.14-2.07)	0.004	1.14 (0.82-1.58)	0.440
Low	1.05 (0.80-1.39)	0.750	1.05 (0.72-1.54)	1.54 (1.17-2.02)	0.002	1.28 (0.89-1.85)	2.93 (2.25-3.81)	< 0.001	1.70 (1.19-2.42)	0.003
VAT mass \geq 75th percentile (ref: < 75th percentile)										
Men										
Adiponectin (tertile)	1.00		1.00	1.00		1.00	1.00		1.00	
High	0.95 (0.58-1.55)	0.826	1.27 (0.74-2.18)	0.86 (0.53-1.41)	0.555	0.95 (0.56-1.59)	1.27 (0.77-2.09)	0.356	1.17 (0.69-2.01)	0.557
Middle	0.98 (0.60-1.61)	0.950	1.39 (0.80-2.41)	0.79 (0.48-1.29)	0.345	0.83 (0.49-1.43)	1.61 (0.99-2.64)	0.056	1.35 (0.79-2.30)	0.274
Low	0.82 (0.51-1.32)	0.414	1.02 (0.54-1.94)	1.79 (1.15-2.78)	0.010	2.29 (1.23-4.25)	2.72 (1.77-4.20)	< 0.001	1.96 (1.08-3.55)	0.027
VAT mass \geq 75th percentile (ref: < 75th percentile)										
Women										
Adiponectin (tertile)	1.00		1.00	1.00		1.00	1.00		1.00	
High	0.91 (0.62-1.34)	0.642	0.99 (0.65-1.49)	0.80 (0.54-1.17)	0.248	0.75 (0.50-1.12)	1.11 (0.75-1.63)	0.607	0.90 (0.59-1.35)	0.603
Middle	1.05 (0.72-1.53)	0.809	1.14 (0.75-1.74)	1.07 (0.73-1.55)	0.737	1.00 (0.66-1.50)	1.50 (1.03-2.18)	0.034	1.06 (0.70-1.60)	0.792
Low	1.20 (0.85-1.71)	0.304	1.07 (0.66-1.72)	1.41 (1.00-1.99)	0.053	0.94 (0.59-1.49)	3.06 (2.19-4.26)	< 0.001	1.59 (1.02-2.49)	0.040
VAT mass \geq 75th percentile (ref: < 75th percentile)										

Values are presented as median (interquartile range) or number (%).

OR = odds ratio, CI = confidence interval, HOMA- β = homeostatic model assessment of β -cell function, VAT = visceral adipose tissue, BMI = body mass index, hs-CRP = high-sensitivity C-reactive protein.

^aAdjusted for gender, age, BMI, hs-CRP, triglyceride, physical activity, adiponectin level, and VAT mass.




Fig. 1. ORs for insulin resistance and β-cell dysfunction according to adiponectin levels and VAT mass. The analyses were adjusted for potential confounders such as gender, age, BMI, hs-CRP, triglyceride, and physical activity.

OR = odds ratio, VAT = visceral adipose tissue, BMI = body mass index, hs-CRP = high-sensitivity C-reactive protein, HOMA-IR = homeostatic model assessment of insulin resistance, HOMA-β = homeostatic model assessment of β-cell function, T1 = tertile 1, T2 = tertile 2, T3 = tertile 3.

^aP < 0.05.

consistent in women (**Supplementary Fig. 1**). When individuals with diabetes were excluded, the combination of a low adiponectin level and high visceral fat mass was associated with the risk of insulin resistance, but not β-cell dysfunction (**Supplementary Tables 1 and 2**).

DISCUSSION

In this cross-sectional study, low adiponectin levels and a high visceral fat mass were associated with an increased risk of insulin resistance and β-cell dysfunction. Obesity is a well-known risk factor for insulin resistance. In particular, abdominal obesity causes insulin resistance by stimulating the formation of metabolic products derived from lipids, hormones, and cytokines. Insulin resistance may lead to endothelial dysfunction and

changes in the insulin signaling pathway.³ Thus, the association between increased visceral fat and the risk of insulin resistance is well-established. In addition, a previous study found an inverse correlation between visceral fat and insulin secretion after adjusting for insulin sensitivity.^{26,27} Further studies reported that visceral fat accumulation contributes to decreased adiponectin levels,^{6,28} although the underlying mechanism is not fully understood. However, some studies suggested that factors that inhibit adiponectin synthesis or secretion are secreted from VAT.²⁹ For example, tumor necrosis factor- α is a strong inhibitor of adiponectin promoter activity.³⁰

Adiponectin mediates systemic insulin sensitivity and glucose homeostasis. A previous study reported that low plasma adiponectin levels predicted abdominal visceral fat accumulation and insulin resistance in Japanese Americans.¹⁵ However, conflicting results on the effects of adiponectin on insulin secretion have been reported so far.^{8,31-34} Some reports suggested a role for adiponectin in modulating plasma insulin levels, both by increasing insulin sensitivity and also via a direct effect on insulin secretion from β -cells. G-adiponectin influences glucose-induced insulin secretion in both human pancreatic islets and rat β -cells.⁸ In a population-based study in Japan, adiponectin levels were inversely associated with HOMA- β .³¹ Although the current study showed that low adiponectin levels were associated with the risk of insulin resistance but not β -cell dysfunction, the combination of low adiponectin levels and a high visceral fat mass was associated with a low disposition index, which is an indicator of β -cell function, relative to insulin sensitivity.

From the result of our study, the effect of adiponectin and abdominal visceral fat on β -cell function was different by gender. Considering the disposition index, the association was significant only in women group. Of course, we cannot rule out the possibility of selection bias, information bias or unexpected confounding factors. However, gender differences in β -cell function should not be underestimated either. First, we have to consider about the difference of adiponectin level and body fat distribution between men and women. A recent research has found that β -cell function has a negative correlation with pancreatic fat mass.³⁵ High visceral fat might increase pancreatic fat and as a result, β -cell function might decrease. Second, there are clinical studies suggesting that women have higher insulin action than men, which supports greater insulin secretion.³⁶ Some studies suggest that gender difference in islet-cell function and susceptibility to failure is a main factor of different insulin action.³⁷ Finally, the different body content or role of sex steroids like estrogen, androgen and progesterone could be also a reason of gender difference in β -cell function.³⁸

We do not know what factors exactly affect the results of our study yet. Further research including prospective study targeting on gender difference is needed.

The current study revealed low adiponectin levels and a high visceral fat mass have combined effects on insulin resistance and β -cell dysfunction. Recent studies revealed a strong association between the combined effects of decreasing adiponectin levels and an increasing visceral fat mass with cardiovascular risk factors.^{9,10} For example, in a population-based study in Japan, a combined effect of an increasing visceral fat mass and decreasing adiponectin level was seen on the prevalence of metabolic syndrome. The ORs for metabolic syndrome with the combination of low adiponectin levels and a high visceral fat mass were 12.7 (95% CI, 9.7-16.6) for men and 13.5 (95% CI, 6.0-30.2) for women.⁹ In Korea, low adiponectin levels and a high visceral fat mass were associated with the development of metabolic syndrome.¹⁰

This study had some limitations that should be addressed. First, due to its cross-sectional design, the direction of causality is unclear, and changes in adiponectin, VAT mass, insulin resistance, and β -cell dysfunction were not monitored. Second, abdominal visceral fat levels were measured by DXA, even though computed tomography is the gold standard for measuring visceral fat.³⁹ Nevertheless, DXA is a valuable tool for assessing visceral fat.^{40,41} Finally, specific cutoff values were used for adiponectin, VAT mass, and glucose metabolism based on a cohort from a specific area in Korea, because there are no population-specific thresholds for those variables.

In conclusion, the current study found that combined low adiponectin levels and a high visceral fat mass might affect not only insulin resistance, but also β -cell dysfunction, even though there could be difference by gender. Examining adiponectin levels in conjunction with visceral fat mass may improve the detection of insulin resistance and β -cell dysfunction. Thus, further prospective studies are needed to explore the combined effects of these factors on the development of insulin resistance and β -cell dysfunction.

SUPPLEMENTARY MATERIALS

Supplementary Table 1

ORs (95% CI) for insulin resistance according to adiponectin levels and VAT mass, after excluding individuals with diabetes

[Click here to view](#)

Supplementary Table 2

ORs (95% CI) for β -cell dysfunction according to adiponectin levels and VAT mass, after excluding individuals with diabetes

[Click here to view](#)

Supplementary Fig. 1

ORs for insulin resistance and β -cell dysfunction according to adiponectin levels and VAT mass by gender. The analyses were adjusted for potential confounders such as age, BMI, hs-CRP, triglyceride, and physical activity. (A) Men. (B) Women.

[Click here to view](#)

REFERENCES

1. Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* 2006;444(7121):840-6.
[PUBMED](#) | [CROSSREF](#)
2. Ibrahim MM. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obes Rev* 2010;11(1):11-8.
[PUBMED](#) | [CROSSREF](#)
3. Balsan GA, Vieira JL, Oliveira AM, Portal VL. Relationship between adiponectin, obesity and insulin resistance. *Rev Assoc Med Bras (1992)* 2015;61(1):72-80.
[PUBMED](#) | [CROSSREF](#)

4. Rabe K, Lehrke M, Parhofer KG, Broedl UC. Adipokines and insulin resistance. *Mol Med* 2008;14(11-12):741-51.
[PUBMED](#) | [CROSSREF](#)
5. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 1999;257(1):79-83.
[PUBMED](#) | [CROSSREF](#)
6. Matsuzawa Y. Establishment of a concept of visceral fat syndrome and discovery of adiponectin. *Proc Jpn Acad Ser B Phys Biol Sci* 2010;86(2):131-41.
[PUBMED](#) | [CROSSREF](#)
7. Kim AY, Park YJ, Pan X, Shin KC, Kwak SH, Bassas AF, et al. Obesity-induced DNA hypermethylation of the adiponectin gene mediates insulin resistance. *Nat Commun* 2015;6(1):7585.
[PUBMED](#) | [CROSSREF](#)
8. Patané G, Caporarello N, Marchetti P, Parrino C, Sudano D, Marselli L, et al. Adiponectin increases glucose-induced insulin secretion through the activation of lipid oxidation. *Acta Diabetol* 2013;50(6):851-7.
[PUBMED](#) | [CROSSREF](#)
9. Matsushita Y, Nakagawa T, Yamamoto S, Kato T, Ouchi T, Kikuchi N, et al. Adiponectin and visceral fat associate with cardiovascular risk factors. *Obesity (Silver Spring)* 2014;22(1):287-91.
[PUBMED](#) | [CROSSREF](#)
10. Cho SA, Joo HJ, Cho JY, Lee SH, Park JH, Hong SJ, et al. Visceral fat area and serum adiponectin level predict the development of metabolic syndrome in a community-based asymptomatic population. *PLoS One* 2017;12(1):e0169289.
[PUBMED](#) | [CROSSREF](#)
11. Okauchi Y, Nishizawa H, Funahashi T, Ogawa T, Noguchi M, Ryo M, et al. Reduction of visceral fat is associated with decrease in the number of metabolic risk factors in Japanese men. *Diabetes Care* 2007;30(9):2392-4.
[PUBMED](#) | [CROSSREF](#)
12. Bi X, Seabolt L, Shibao C, Buchowski M, Kang H, Keil CD, et al. DXA-measured visceral adipose tissue predicts impaired glucose tolerance and metabolic syndrome in obese Caucasian and African-American women. *Eur J Clin Nutr* 2015;69(3):329-36.
[PUBMED](#) | [CROSSREF](#)
13. Inoue M, Maehata E, Yano M, Taniyama M, Suzuki S. Correlation between the adiponectin-leptin ratio and parameters of insulin resistance in patients with type 2 diabetes. *Metabolism* 2005;54(3):281-6.
[PUBMED](#) | [CROSSREF](#)
14. Weber KS, Strassburger K, Pacini G, Nowotny B, Müsigg K, Szendroedi J, et al. Circulating adiponectin concentration is inversely associated with glucose tolerance and insulin secretion in people with newly diagnosed diabetes. *Diabet Med* 2017;34(2):239-44.
[PUBMED](#) | [CROSSREF](#)
15. Han SJ, Boyko EJ, Fujimoto WY, Kahn SE, Leonetti DL. Low plasma adiponectin concentrations predict increases in visceral adiposity and insulin resistance. *J Clin Endocrinol Metab* 2017;102(12):4626-33.
[PUBMED](#) | [CROSSREF](#)
16. Oka R, Yagi K, Sakurai M, Nakamura K, Nagasawa SY, Miyamoto S, et al. Impact of visceral adipose tissue and subcutaneous adipose tissue on insulin resistance in middle-aged Japanese. *J Atheroscler Thromb* 2012;19(9):814-22.
[PUBMED](#) | [CROSSREF](#)
17. Scheuer SH, Færch K, Philipsen A, Jørgensen ME, Johansen NB, Carstensen B, et al. Abdominal fat distribution and cardiovascular risk in men and women with different levels of glucose tolerance. *J Clin Endocrinol Metab* 2015;100(9):3340-7.
[PUBMED](#) | [CROSSREF](#)
18. Shim JS, Song BM, Lee JH, Lee SW, Park JH, Choi DP, et al. Cardiovascular and Metabolic Diseases Etiology Research Center (CMERC) cohort: study protocol and results of the first 3 years of enrollment. *Epidemiol Health* 2017;39:e2017016.
[PUBMED](#) | [CROSSREF](#)
19. Kaul S, Rothney MP, Peters DM, Wacker WK, Davis CE, Shapiro MD, et al. Dual-energy X-ray absorptiometry for quantification of visceral fat. *Obesity (Silver Spring)* 2012;20(6):1313-8.
[PUBMED](#) | [CROSSREF](#)
20. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28(7):412-9.
[PUBMED](#) | [CROSSREF](#)
21. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. *Diabetes Care* 1999;22(9):1462-70.
[PUBMED](#) | [CROSSREF](#)

22. Bergman RN, Belfiore F, Molinatti GM. Current topics in diabetes research. In: Turner R, Levy J, Rudenski A, Hammersley M, Page R, editors. *Measurement of Insulin Resistance and Beta-cell Function: the HOMA and CIGMA Approach*. Berlin: Karger Publishers; 1993, 66-75.
23. Goedecke JH, Dave JA, Faulenbach MV, Utzschneider KM, Lambert EV, West S, et al. Insulin response in relation to insulin sensitivity: an appropriate beta-cell response in black South African women. *Diabetes Care* 2009;32(5):860-5.
[PUBMED](#) | [CROSSREF](#)
24. Utzschneider KM, Prigeon RL, Faulenbach MV, Tong J, Carr DB, Boyko EJ, et al. Oral disposition index predicts the development of future diabetes above and beyond fasting and 2-h glucose levels. *Diabetes Care* 2009;32(2):335-41.
[PUBMED](#) | [CROSSREF](#)
25. Yamada C, Mitsuhashi T, Hiratsuka N, Inabe F, Araidai N, Takahashi E. Optimal reference interval for homeostasis model assessment of insulin resistance in a Japanese population. *J Diabetes Investig* 2011;2(5):373-6.
[PUBMED](#) | [CROSSREF](#)
26. Utzschneider KM, Carr DB, Hull RL, Kodama K, Shofer JB, Retzlaff BM, et al. Impact of intra-abdominal fat and age on insulin sensitivity and beta-cell function. *Diabetes* 2004;53(11):2867-72.
[PUBMED](#) | [CROSSREF](#)
27. Wagenknecht LE, Langefeld CD, Scherzinger AL, Norris JM, Haffner SM, Saad MF, et al. Insulin sensitivity, insulin secretion, and abdominal fat: the Insulin Resistance Atherosclerosis Study (IRAS) Family Study. *Diabetes* 2003;52(10):2490-6.
[PUBMED](#) | [CROSSREF](#)
28. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. 1999. *Biochem Biophys Res Commun* 2012;425(3):560-4.
[PUBMED](#) | [CROSSREF](#)
29. Halleux CM, Takahashi M, Delporte ML, Detry R, Funahashi T, Matsuzawa Y, et al. Secretion of adiponectin and regulation of apM1 gene expression in human visceral adipose tissue. *Biochem Biophys Res Commun* 2001;288(5):1102-7.
[PUBMED](#) | [CROSSREF](#)
30. Maeda N, Takahashi M, Funahashi T, Kihara S, Nishizawa H, Kishida K, et al. PPARgamma ligands increase expression and plasma concentrations of adiponectin, an adipose-derived protein. *Diabetes* 2001;50(9):2094-9.
[PUBMED](#) | [CROSSREF](#)
31. Nakamura A, Miyoshi H, Ukawa S, Nakamura K, Nakagawa T, Terauchi Y, et al. Serum adiponectin and insulin secretion: a direct or inverse association? *J Diabetes Investig* 2018;9(5):1106-9.
[PUBMED](#) | [CROSSREF](#)
32. Winzell MS, Nogueiras R, Dieguez C, Ahrén B. Dual action of adiponectin on insulin secretion in insulin-resistant mice. *Biochem Biophys Res Commun* 2004;321(1):154-60.
[PUBMED](#) | [CROSSREF](#)
33. Staiger K, Stefan N, Staiger H, Brendel MD, Brandhorst D, Bretzel RG, et al. Adiponectin is functionally active in human islets but does not affect insulin secretory function or beta-cell lipooptosis. *J Clin Endocrinol Metab* 2005;90(12):6707-13.
[PUBMED](#) | [CROSSREF](#)
34. Okamoto M, Ohara-Imaizumi M, Kubota N, Hashimoto S, Eto K, Kanno T, et al. Adiponectin induces insulin secretion in vitro and in vivo at a low glucose concentration. *Diabetologia* 2008;51(5):827-35.
[PUBMED](#) | [CROSSREF](#)
35. Roh E, Kim KM, Park KS, Kim YJ, Chun EJ, Choi SH, et al. Comparison of pancreatic volume and fat amount linked with glucose homeostasis between healthy Caucasians and Koreans. *Diabetes Obes Metab* 2018;20(11):2642-52.
[PUBMED](#) | [CROSSREF](#)
36. Basu R, Dalla Man C, Campioni M, Basu A, Klee G, Toffolo G, et al. Effects of age and sex on postprandial glucose metabolism: differences in glucose turnover, insulin secretion, insulin action, and hepatic insulin extraction. *Diabetes* 2006;55(7):2001-14.
[PUBMED](#) | [CROSSREF](#)
37. Gannon M, Kulkarni RN, Tse HM, Mauvais-Jarvis F. Sex differences underlying pancreatic islet biology and its dysfunction. *Mol Metab* 2018;15(9):82-91.
[PUBMED](#) | [CROSSREF](#)
38. Mauvais-Jarvis F. Role of sex steroids in β cell function, growth, and survival. *Trends Endocrinol Metab* 2016;27(12):844-55.
[PUBMED](#) | [CROSSREF](#)

39. Rössner S, Bo WJ, Hiltbrandt E, Hinson W, Karstaedt N, Santago P, et al. Adipose tissue determinations in cadavers--a comparison between cross-sectional planimetry and computed tomography. *Int J Obes* 1990;14(10):893-902.
[PUBMED](#)
40. Jung SH, Ha KH, Kim DJ. Visceral fat mass has stronger associations with diabetes and prediabetes than other anthropometric obesity indicators among Korean adults. *Yonsei Med J* 2016;57(3):674-80.
[PUBMED](#) | [CROSSREF](#)
41. Snijder MB, Visser M, Dekker JM, Seidell JC, Fuerst T, Tylavsky F, et al. The prediction of visceral fat by dual-energy X-ray absorptiometry in the elderly: a comparison with computed tomography and anthropometry. *Int J Obes Relat Metab Disord* 2002;26(7):984-93.
[PUBMED](#) | [CROSSREF](#)