

## Corrigendum: Dental alloplastic bone substitutes currently available in Korea

Jeong-Kui Ku<sup>1,2</sup>, Inseok Hong<sup>3</sup>, Bu-Kyu Lee<sup>1</sup>, Pil-Young Yun<sup>4</sup>, Jeong Keun Lee<sup>5</sup>

<sup>1</sup>Department of Oral and Maxillofacial Surgery, Asan Medical Center, Seoul, <sup>2</sup>Department of Oral and Maxillofacial Surgery, Section of Dentistry, Armed Forces Capital Hospital, Seongnam, <sup>3</sup>Department of Oral and Maxillofacial Surgery, School of Dentistry and Institute of Oral Bioscience, Research institute of Clinical Medicine of Chonbuk National University-Biomedical Research Institute of Chonbuk National University Hospital, Chonbuk National University, Jeonju, <sup>4</sup>Department of Oral and Maxillofacial Surgery, Section of Dentistry, Seoul National University Bundang Hospital, Seongnam,

<sup>5</sup>Department of Oral and Maxillofacial Surgery, Institute of Oral Health Science, Ajou University School of Medicine, Suwon, Korea

J Korean Assoc Oral Maxillofac Surg 2019;45:51-67.

https://doi.org/10.5125/jkaoms.2019.45.2.51

This correction is being published to correct the 4th author's affiliation in above article. The authors apologize for this error.

Before correction

Jeong-Kui Ku<sup>1,2</sup>, Inseok Hong<sup>3</sup>, Bu-Kyu Lee<sup>1</sup>, Pil-Young Yun<sup>2</sup>, Jeong Keun Lee<sup>4</sup>

<sup>1</sup>Department of Oral and Maxillofacial Surgery, Asan Medical Center, Seoul, <sup>2</sup>Department of Oral and Maxillofacial Surgery, Section of Dentistry, Armed Forces Capital Hospital, Seongnam, <sup>3</sup>Department of Oral and Maxillofacial Surgery, School of Dentistry and Institute of Oral Bioscience, Research Institute of Clinical Medicine of Chonbuk National University-Biomedical Research Institute of Chonbuk National University Hospital, Chonbuk National University, Jeonju, <sup>4</sup>Department of Oral and Maxillofacial Surgery, Institute of Oral Health Science, Ajou University School of Medicine, Suwon, Korea

After correction

Jeong-Kui Ku<sup>1,2</sup>, Inseok Hong<sup>3</sup>, Bu-Kyu Lee<sup>1</sup>, Pil-Young Yun<sup>4</sup>, Jeong Keun Lee<sup>5</sup>

<sup>1</sup>Department of Oral and Maxillofacial Surgery, Asan Medical Center, Seoul,

<sup>2</sup>Department of Oral and Maxillofacial Surgery, Section of Dentistry, Armed Forces Capital Hospital, Seongnam, <sup>3</sup>Department of Oral and Maxillofacial Surgery, School of Dentistry and Institute of Oral Bioscience, Research institute of Clinical Medicine of Chonbuk National University-Biomedical Research Institute of Chonbuk National University Hospital, Chonbuk National University, Jeonju, <sup>4</sup>Department of Oral and Maxillofacial Surgery, Section of Dentistry, Seoul National University Bundang Hospital, Seongnam, <sup>5</sup>Department of Oral and Maxillofacial Surgery, Institute of Oral Health Science, Ajou University School of Medicine, Suwon, Korea

(© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/4.0/), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Copyright © 2019 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.