

Original Article
Medicine General & Policy



Systematic Preventable Trauma Death Rate Survey to Establish the Region-based Inclusive Trauma System in a Representative Province of Korea

Junsik Kwon ,^{1,2} Jin-Hee Lee ,³ Kyungjin Hwang ,^{1,2} Yunjung Heo ,⁴
Hang Joo Cho ,⁵ John Cook-Jong Lee ,^{1,2} and Kyoungwon Jung ^{1,2}

OPEN ACCESS

Received: Jun 17, 2020
Accepted: Oct 15, 2020

Address for Correspondence:

Kyoungwon Jung, MD, PhD

Division of Trauma Surgery, Department of Surgery, Ajou University School of Medicine, 164 World cup-ro, Yeongtong-gu, Suwon 16499, Republic of Korea.
E-mail: jake98@daum.net

© 2020 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Junsik Kwon
<https://orcid.org/0000-0003-3164-4995>
Jin-Hee Lee
<https://orcid.org/0000-0002-2556-5484>
Kyungjin Hwang
<https://orcid.org/0000-0002-7664-8503>
Yunjung Heo
<https://orcid.org/0000-0001-5708-1428>
Hang Joo Cho
<https://orcid.org/0000-0003-3322-744X>
John Cook-Jong Lee
<https://orcid.org/0000-0002-5468-3340>
Kyoungwon Jung
<https://orcid.org/0000-0001-7895-0362>

Funding

This study was supported by the Local Government of Gyeonggi Province in Korea in 2018, through the project for the establishment of a regional trauma system.

<https://jkms.org>

¹Division of Trauma Surgery, Department of Surgery, Ajou University School of Medicine, Suwon, Korea
²Southern Gyeonggi Trauma Center, Ajou University Hospital, Suwon, Korea
³National Emergency Medical Center, National Medical Center, Seoul, Korea
⁴The Health Insurance Review & Assessment Service, Wonju, Korea
⁵Department of Emergency Medicine, Uijeongbu St. Mary's Hospital, The Catholic University of Korea, Seoul, Korea

ABSTRACT

Background: Trauma mortality review is the first step in assessing the quality of the trauma treatment system and provides an important basis for establishing a regional inclusive trauma system. This study aimed to obtain a reliable measure of the preventable trauma death rate in a single province in Korea.

Methods: From January to December 2017, a total of 500 sample cases of trauma-related deaths from 64 hospitals in Gyeonggi Province were included. All cases were evaluated for preventability and opportunities for improvement using a multidisciplinary panel review approach.

Results: Overall, 337 cases were included in the calculation for the preventable trauma death rate. The preventable trauma death rate was estimated at 17.0%. The odds ratio was 3.97 folds higher for those who arrived within “1–3 hours” than those who arrived within “1 hour.” When the final treatment institution was not a regional trauma center, the odds ratio was 2.39 folds higher than that of a regional trauma center. The most significant stage of preventable trauma death was the hospital stage, during which 86.7% of the cases occurred, of which only 10.3% occurred in the regional trauma center, whereas preventable trauma death was more of a problem at emergency medical institutions.

Conclusion: The preventable trauma death rate was slightly lower in this study than in previous studies, although several problems were noted during inter-hospital transfer; in the hospital stage, more problems were noted at emergency medical care facilities than at regional trauma centers. Further, several opportunities for improvements were discovered regarding bleeding control.

Keywords: Injuries and Wounds; Trauma System; Trauma Center; Preventable Death Rate; Province; Korea

INTRODUCTION

Trauma is the leading cause of death worldwide, in adults aged ≤ 39 years, and is one of the greatest health problems in both low and high-income countries.¹ However, systematic efforts to lower

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Kwon J, Lee JC, Hwang K, Heo Y, Jung K. Data curation: Kwon J, Lee JH, Hwang K. Formal analysis: Kwon J, Lee JH, Hwang K, Jung K. Investigation: Kwon J, Hwang K, Cho HJ, Jung K. Methodology: Kwon J, Lee JC, Hwang K, Heo Y, Lee JCJ, Jung K. Writing - original draft: Kwon J, Lee JH, Hwang K, Jung K. Writing - review & editing: Kwon J, Lee JC, Hwang K, Heo Y, Cho HJ, Lee JCJ, Jung K.




Fig. 1. Emergency medical facilities including regional trauma centers in Gyeonggi Province.

trauma-related deaths are limited; with the exception of a few countries with well-established trauma systems, no countries have shown a marked reduction in trauma-related death rates.² Most healthcare fields in Korea have made some progress; however, trauma care remains at a rudimentary stage, and continuous quality control using consistent and objective scales is lacking.

Preventable trauma death review is a classic quality control trauma care systems tool, defined as the review of trauma-related avoidable deaths in which the patient would have survived if transferred to an appropriate hospital in time for appropriate treatment.³ Despite limitations, this index is still important in determining the performance of trauma care systems. The preventable trauma death rate has been measured inconsistently using different methods in Korea, with reported rates of 50.4% in the late 1990s, 35.2% in 2009–2010, and 30.5% in 2015.⁴⁻⁶

Gyeonggi-do, the region evaluated in our study, is about the size of the Los Angeles County in the United States. It is densely populated with approximately 13.09 million people, a quarter of the entire Korean population.⁷ Along with Seoul, Gyeonggi-do is the most urbanized and medically dense Korean region with 57 emergency medical care facilities, including five regional emergency medical centers at the time of survey (**Fig. 1**). However, compared with the high accessibility to emergency medical care, its trauma care system, similar to many others in Korea, is still at a rudimentary stage with multiple problems. A review of trauma-related deaths in this region can be used as the basis for reorganizing resources and establishing inclusive trauma systems in the region.

This study aimed to obtain a reliable measure of the preventable trauma death rate in a single province as of 2017, analyze the related problems, and obtain foundational data for devising enhancement measures that can be used for regional-level quality control, to advance the national trauma system in Korea.

METHODS

Study population

This study included data on those who experienced trauma-related deaths from 64 hospitals providing emergency medical care in the Gyeonggi Province from January 1, 2017 to December 31, 2017. Trauma-related death was defined as the deaths of patients who had presented to the emergency department (ED) and had one or more S or T codes as per the Korean Standard Classification of Diseases (KCD) seventh edition (nearly identical to the 10th revision of the International Statistical Classification of Diseases and Related Health Problems) in their diagnosis while exiting the ED or hospital (excluding patients with a code for frostbite, addiction, intoxication, drowning, burns, other and unspecified effects of external causes, or complications regarding surgical and medical care). As of December 31, 2017, 400 out of 402 total emergency medical care facilities in Korea had registered their data (99.5%); 64 emergency medical care facilities in Gyeonggi Province were registered. Overall, 1,546 trauma-related deaths were registered by 57 of the 64 hospitals in Gyeonggi Province. Of these, 263 patients were dead on arrival (DOA) (17.0%), 97 deaths occurred in the ED (6.3%), 848 deaths occurred after hospital admission (54.8%), and 338 deaths were after transfer (21.9%).

Sampling

The study and target populations were identical in background characteristics, and to compute a regionally representative preventable trauma death rate in Gyeonggi region, a stratified two-stage cluster sampling method was applied, and stratification was designed as a double layer (Fig. 2). The National Emergency Department Information System (NEDIS), a database containing information on patients presenting to the ED in Korea, with documentation of the treatment process in real-time, was used as the sampling frame. Sampling was performed on all cases in facilities with deaths after transfer to a second facility, according to the sample size of facilities with no deaths after transfer to a second




Fig. 2. Study design and sampling.
NEDIS = National Emergency Department Information System.

facility. Sample size for each facility was set according to the type of hospital (regional trauma center [RTC], regional emergency medical center, local emergency medical center, local emergency medical service), number of deaths (≥ 100 , 50–99, 30–49, 10–29, < 10), place of death (DOA in first facility, death in ED in the first facility, death after hospital admission in the first facility, death after transfer to second facility), and patient age (≤ 14 years, 15–54 years, ≥ 55 years). For patients aged ≤ 14 years, the entire patient pool was sampled. Deaths from all RTCs were sampled, and deaths from all regional emergency medical centers, except for one center, were sampled. The extracted sample consisted of 478 cases from 42 facilities, with deaths after transfer to a second facility, and 22 cases from six facilities without deaths after transfer to a second facility, totaling 500 cases from 48 facilities (80 cases from 2 RTCs, 75 cases from four regional emergency medical centers, 265 cases from 23 local emergency medical centers, and 80 cases from 19 local emergency medical services).

Data collection and type of data

With permission from the local government, patient medical records were collected from hospitals that transferred the sampled trauma patients to the 48 study hospitals. Based on the preventable trauma death review form (**Supplementary Data 1**) that we developed, the study hospitals submitted the necessary data and completed checklists. Types of data included emergency care record, death certificate, CD containing image copies, and copy of medical record. The copy of the medical record had to contain initial care record at the ED, the progress note, nursing care record, imaging and blood test results, and admission/discharge record. We requested additional data when needed, and medical records were collected from the 48 hospitals in which the trauma patients had died and from the 221 hospitals that had transferred these patients to the final hospitals to investigate the 500 cases included in our study.

Composition of designated assistants and preventable death panel and multidisciplinary reviewing

The preventable trauma death review form was based on the preventable trauma death evaluation form with an embedded audit filter as in the World Health Organization (WHO) guidelines for the trauma quality improvement program.¹ Designated assistants consisting of seven trauma coordinators working in RTCs nationwide investigated and recorded the general characteristics of patients, injury-related information, and transport and treatment-related information using the preventable trauma death review form prior to the review by the preventable death panel. The preventable death panel consisting of 15 dedicated trauma professionals working in RTCs nationwide, performed an initial evaluation of the preventability of deaths using the data extracted by the designated assistants. For a multidisciplinary review, three teams were formed, with each team consisting of two general surgeons, one thoracic surgeon, one neurosurgeon, and one emergency physician. After individual evaluations, the panels from each team came together to present the cases they had reviewed, and they discussed and reached a consensus for a final assessment of the preventability of trauma-related deaths.

Criteria for preventable trauma deaths

The criteria presented by the WHO guidelines for trauma quality improvement programs¹ were used as the criteria for the preventability of trauma-related deaths. These criteria assess preventability in consideration of the severity of trauma and appropriateness of the provided treatment. Preventable trauma deaths were computed by summing “preventable (P)” and “potentially preventable (PP)” deaths.

Qualitative analysis

The preventable trauma death rate was computed using the selected sample; we also extracted data on patients who died after inter-hospital transfer, who showed a substantially high preventable trauma death rate in a previous study,⁴ and patients who died after hospital admission. These patients having the highest percentage of various trauma death stages were therefore analyzed to provide feedback to enhance the quality of trauma patient care. These additional data were only used for qualitative analysis and were not included in the computation of the preventable trauma death rate. All preventable trauma deaths included in the qualitative analysis were classified according to five causes of death: bleeding, sepsis/multi-organ dysfunction syndrome (MODS), central nervous system (CNS) injury, respiratory failure, and cardiac issues. One main cause of death was chosen, but more than one cause of death could be recorded.

Statistical analyses

The population was estimated using the weighted values of sampling rates. The methods used for mortality estimation and adjustment in a previous study⁴ were used. Risk factors of preventable trauma deaths were identified using multiple logistic regression analysis, and variables previously identified to potentially impact the preventable trauma death rate were used as explanatory variables. Variables included in the logistic regression model were age, type of hospital, inter-hospital transfer, time from accident to death, time from accident to arrival at the hospital, and mechanism of injury. Risk factors were analyzed using the observed number of death cases instead of the estimated population. We also analyzed consistency among panels using the Cohen's Kappa index to establish the reliability of the results.

Ethics statement

This study was approved by the Institutional Review Board of Ajou University (AJIRB-MED-EXP-18-157). The requirement for informed consent was waived by the board due to the observational nature of the study.

RESULTS

General characteristics

After excluding 163 patients (four patients, death was not clearly confirmed; 151 patients, cause of death was non-traumatic; and eight patients, insufficient data) from the total sample of 500 patients, the preventable trauma death rate was computed based on 337 patients. **Table 1** shows the baseline characteristics of the subjects. In addition to the values for stratified samples, estimates calculated based on them and percentage in the population are also presented in **Table 1**. There were more men in our sample (67.1%), with a mean age of 59 years, and 75.7% of the sample was ≥ 55 years. The most common type of hospital in which trauma-related deaths occurred was the local emergency center (48.4%), and approximately 77.2% of patients presented to the center via the 119 emergency medical service (EMS) instead of being transferred from another hospital. The most common mechanism of injury was a traffic accident (34.7%), and in most cases, the time from accident to death was ≥ 7 days (27.7%). In most cases, the time from accident to arrival at the hospital was < 1 hour (51.8%).

Preventable trauma death rate and relevant factors

The preventable trauma death rate in Gyeonggi region in 2017 was estimated to be 17.0% (95% confidence interval [CI], 13.7–20.3), and 2.9% (95% CI, 0.0–6.2) of these deaths were

Table 1. Baseline characteristics of the study participants (n = 337) (estimated n = 1,546)

Characteristics	Patients, No.	Estimated patients, No.	Weighted %
Sex			
Male	225	1,037	67.1
Female	112	509	32.9
Age, yr			
≤ 14	20	23	1.5
15–54	83	352	22.8
≥ 55	234	1,171	75.7
Type of emergency care center			
Regional trauma center	74	368	23.8
Regional emergency center	52	237	15.3
Local emergency center	169	748	48.4
Local emergency service	42	193	12.5
Transfer, y/n			
No, directly visited the last hospital	261	1,194	77.2
Yes, once via another hospital	69	323	20.9
Yes, more than two folds via another hospital	7	29	1.8
Mode of transport			
119 + air-transport	257	1,176	76.1
129 + hospital ambulance	61	282	18.2
Others	19	88	5.7
Mechanism of injury			
Motor vehicle crash	125	536	34.7
Fall from a height	76	350	22.6
Ground-level falls	75	361	23.4
Others	25	122	7.9
Unknown	36	177	11.4
Time from accident to death			
Time < 1 hr	50	194	12.6
1 hr ≤ time < 6 hr	51	211	13.6
6 hr ≤ time < 24 hr	26	135	8.7
1 day ≤ time < 2 day	20	94	6.1
2 day ≤ time < 7 day	61	304	19.7
7 day ≤ time	90	429	27.7
Unknown	39	179	11.6
Time from accident to arrival at the hospital			
Time < 1 hr	179	800	51.8
1 hr ≤ time < 3 hr	50	245	15.9
3 hr ≤ time < 6 hr	31	152	9.8
6 hr ≤ time < 1 day	35	155	10.0
Unknown	42	194	12.5

P deaths, while 14.1% (95% CI, 11.1–17.1) were PP deaths (Table 2). The Cohen's Kappa index of inter-rater reliability, measured after randomizing and sampling 120 cases, was 0.25 (fair reliability). The time from accident to arrival at the hospital influenced the preventable trauma death rate, though the risk was higher in patients who arrived at the ED within 1–3 hours ($P = 0.01$) and those who arrived within 3–24 hours ($P = 0.09$) of the accident than in patients who arrived within 1 hour of the accident. No significant differences were observed despite the risk being higher in patients who died < 6 hours after the accident ($P = 0.06$) compared with those who died between 6–24 hours after the accident. The risk was also higher in patients admitted to non-RTC emergency care centers than in patients admitted to RTCs ($P = 0.06$) (Table 3).

Percentage of impact on preventable trauma death rate by stage and causes

Stages that affected preventable trauma deaths occurring in emergency care centers in Gyeonggi Province were classified into the pre-hospital stage, inter-hospital transfer

Table 2. Preventable trauma death rate (estimated n = 1,546)

Variables	Estimated patients, No. (%)	95% CI
PTD (P+PP)	263 (17.0)	13.7–20.3
P	44 (2.9)	0.0–6.2
PP	219 (14.1)	11.1–17.1
Non-PTD (NP+NPCI)	1,283 (83.0)	79.6–86.4
NP	937 (60.6)	57.5–63.8
NPCI	346 (22.4)	18.6–26.2

CI = confidence interval, PTD = preventable trauma death, P = preventable, PP = potentially preventable, NP = non-preventable, NPCI = non-preventable, but with care that could have been improved.

Table 3. Factors associated with preventable trauma death (n = 337)

Variables	P+PP	NP+NPCI	Odds ratio	P value
Age, yr				
≤ 14	2	18	1	
15–54	9	74	1.01	0.99
≥ 55	44	190	1.40	0.68
Type of hospital				
Regional trauma center	8	66	1	
Others	47	216	2.39	0.06
Transfer, y/n				
Directly visited	38	223	1	
Transferred	17	59	1.08	0.85
Time from accident to death				
Time < 6 hr	10	91	1	
6 hr ≤ time < 24 hr	7	19	2.99	0.07
24 hr ≤ time < 7 day	16	65	1.99	0.19
7 day ≤ time	8	82	0.60	0.40
Time from accident to arrival at the hospital				
Time < 1 hr	17	162	1	
1 hr ≤ time < 3 hr	14	36	3.97	0.01
3 hr ≤ time < 24 hr	10	56	2.55	0.09
Mechanism of injury				
Motor vehicle crash	23	102	1	
Fall from a height	11	65	0.50	0.14
Ground-level falls	15	60	0.46	0.11
Others	4	21	0.41	0.19

P = preventable, PP = potentially preventable, NP = non-preventable, NPCI = non-preventable, but with care that could have been improved.

stage, and hospital stage. The hospital stage (86.7%) had the greatest impact, followed by inter-hospital transfer stage (8.4%) and pre-hospital stage (4.9%). Of the 86.7% patients experiencing problems at the hospital stage, only 10.3% experienced them in an RTC, while 76.4% experienced them in a hospital other than an RTC (Table 4). The causes of death in preventable cases included hemorrhage in 39 (42.9%) cases, sepsis/MODS in 26 (28.6%) cases, CNS injury in 14 (15.4%) cases, respiratory failure in 11 (12.1%) cases, and cardiac failure in two (2.2%) cases. In one patient who experienced hemorrhage, the contribution to “death from traumatic brain injury and hemorrhagic shock” was considered to be identical and duplicated.

Qualitative analysis of preventable trauma deaths

In addition to the 337 effective samples selected for determining the preventable trauma death rate, we analyzed an additional 10% of cases classified under “deaths after inter-hospital transfer” and “deaths after admission,” which were subgroups with the highest preventable trauma death rates in a previous study⁴; in total, 524 patients were included in the analysis of trauma-related death. Of the 524 patients, 93 were determined to have experienced preventable trauma-related deaths. The most common cause of death in

Table 4. Location of deficit and hospital types according to the preventable trauma death rate (n = 263)

Location of deficiencies	Preventable trauma death		
	Estimated patients, No.	Preventable trauma death rate	%
Total	263	17.0	100.0
Pre-hospital	13	0.8	4.9
Inter-hospital	22	1.5	8.4
Hospital	228	14.7	86.7
Regional trauma center	34	2.2	12.9
Inter-hospital	7	0.5	2.7
Hospital	27	1.7	10.3
Emergency care centers other than regional trauma centers	229	14.8	87.1
Pre-hospital	13	0.8	4.9
Inter-hospital	15	1.0	5.7
Hospital	201	13.0	76.4

preventable trauma-related deaths was hemorrhage, followed by sepsis/MODS, CNS injury, respiratory failure, and cardiac issues (Table 5). The opportunities for improvement (OFIs) in 39 patients in whom hemorrhage was the cause of death were analyzed and presented in Table 6, with a duplicate selection being included. Among the hospital-level problems, delayed surgery or embolization for hemostasis, experienced by 30 patients, was the most common OFI, followed by delayed/insufficient transfusion (experienced by 25 patients).

Table 5. Causes of death in preventable cases (n = 93)

Causes of death	Cases, No. (%)
Hemorrhage ^a	39 (42.9)
Sepsis/MODS	26 (28.6)
CNS injury ^a	14 (15.4)
Respiratory failure	11 (12.1)
Cardiac issue	2 (2.2)

MODS = multi-organ dysfunction syndrome, CNS = central nervous system.

^aIn one case, the contribution of hemorrhage and CNS injury to the cause of death were considered identical and duplicated.

Table 6. Opportunities for improvement in preventable trauma deaths according to main cause of death and location of deficiency

Main cause of death ^a	Location of deficiencies	Opportunities for improvement	Cases, No. ^b
Hemorrhage	Pre-hospital	Failure to achieve vascular access	6
		Field triage failure	5
		Delayed on-site treatment	1
	Inter-hospital transfer	Delayed transportation	9
		Problem with appropriate airway management	3
		Delayed transfusion	2
		Inappropriate transfer	1
	Hospital	Delayed surgery/embolization for hemostasis	30
		Delayed/insufficient transfusion	25
		Delay in other treatments	11
		Problem with airway management and ventilation	8
		Insufficient records	6
		Dangerous CT	5
Sepsis/MODS	Hospital	Shortage of ORs or ICUs	2
		Problem related to caregivers	1
CNS injury	Hospital	Other	1
		Delayed diagnosis/treatment of cause of sepsis	26
CNS injury	Hospital	Delayed craniotomy	8
		Failure to perform craniotomy	6

CT = computed tomography, OR = operation room, ICU = intensive care unit, MODS = multi-organ dysfunction syndrome, CNS = central nerve system.

^aThe three most common causes of death are presented. ^bA single case may appear multiple times due to two or more identified opportunities for improvement.

We observed delays in diagnosis and intensive care treatment in 26 patients who died from sepsis/MODS. All 14 patients who died from CNS injuries had traumatic brain injury, with delayed craniotomy or failure to perform craniotomy as the main OFIs in these cases. When craniotomy was performed, the average time from arrival at the final destined hospital to surgery was 8.7 hours (median value, inter-quartile range; 3.3–15.0 hours).

DISCUSSION

The first reported preventable trauma death rate in Korea was 50.4% in 1997⁸; since then, there has been a consistent decline in this rate, with the most recent report showing a rate of 30.5% in 2015, suggesting an improvement.⁴ This rate is similar to that reported by Motomura et al.⁹ in Chiba Prefecture in 2011 (29.0%), although it is still higher than those reported in countries with stable trauma care systems, such as the United States and Canada. In 2005 in the US, the rate was 6.7% in Utah,¹⁰ and 2.4% in California, in 1998.¹¹ The preventable trauma death rate in Gyeonggi Province in 2017 was 17.0% according to this study, which is a substantial improvement from that reported in a national survey using 2015 data.⁴ This may be attributable to the high accessibility to trauma care in Gyeonggi Province, as it is a medically dense area compared with other regions in Korea. The establishment of diverse emergency medical care facilities that have not been designed for trauma care may poorly reflect the actual outcome of the trauma care system.¹² If the improvement in the preventable trauma death rate in Gyeonggi region found in this study is a result of enhancement in the quality of the trauma care system, this could be attributable to the two RTCs in southern/northern Gyeonggi Province that opened in 2016 and 2017, respectively.

The time from accident to arrival at the hospital and RTC as the final care institution were identified to have significant effects on the determination of preventability. This is understandable since reducing the time taken from the accident to definitive treatment and establishing designated trauma centers are critical to the success of early trauma care systems worldwide. However, delayed transfer to the final treatment facility seems to be affected by the transfer process instead of the initial transportation in the pre-hospital stage. Unnecessary inter-hospital transfers and excessive testing and procedures performed at transferring hospitals resulting in delays in patient arrival at the final treatment facility have been reported in a prior study conducted in Gyeonggi Province.⁸ Twenty-seven of the 228 patients who experienced preventable trauma deaths due to problems in the hospital stage were speculated to have died due to problems at an RTC (Table 4). Considering that 23.8% of all trauma-related deaths (368/1,546) occurred in an RTC, the number of trauma-related deaths that had occurred in one of the 62 hospitals other than an RTC in Gyeonggi Province was more than three-fold higher than that in one of the two RTCs (368 vs. 1,178); however, there were 7.4 folds more hospital-stage problems that led to preventable trauma deaths (27 vs. 201). This highlights the distinct difference in the capacity to treat severely injured patients, between RTCs and other hospitals. Assuming that RTCs have an unlimited pool of resources and thus can accept an unlimited number of patients, inter-hospital transfer complications could have been prevented if patients were transferred to an RTC for initial treatment. In such cases, we can substitute the fraction of preventable trauma deaths that occurred in the hospital stage in hospitals other than an RTC (13%) with the fraction of preventable trauma deaths that occurred in an RTC (1.7%); this leads to a total preventable trauma death rate of 4.2% (pre-hospital stage 0.8% + inter-hospital transfer stage 0% + hospital stage 3.4% [1.7%+1.7%]). With this calculation, we can estimate that 198 regional

trauma patients could be additionally saved per year. With a total preventable trauma death estimate of 263, preventable trauma deaths can be lowered by 75.3% (198/263) according to our findings. However, trauma-related deaths were calculated based on the NEDIS database; a higher estimate of trauma-related deaths may be derived if the trauma-related deaths that were not registered in the NEDIS were taken into consideration. Although this is a simple arithmetic estimate that disregards the capacity of RTCs for accepting patients, it presents implications for the direction of future regional trauma systems. Currently, RTCs in southern and northern Gyeonggi together have 173 bed-spaces, and they serve a function similar to that of a level 1 trauma center, as defined in the resources for optimal care of the injured patient.¹³ Considering the 12-million population in the region and the lack of designated trauma care facilities other than a level 1 facility, the capacities of these RTCs are substantially low. This can be addressed by establishing a truly inclusive regional trauma system, by restructuring the available medical resources, making them trauma care appropriate, and by modifying the pre-hospital transport system so that severe trauma patients are directly transported to an RTC. Thus, a consultative organization that can coordinate non-RTC emergency medical care facilities, 119 EMS, and local governments with the existing RTCs is needed, and the recently founded Gyeonggi Trauma Systems Organization could play an important role to this end.

Instead of merely calculating the preventable trauma death rate, this study ultimately aimed to enhance the quality of trauma care by providing constructive feedback to facilities that encountered preventable trauma deaths. For this purpose, we performed a qualitative analysis by including an additional 10% of “deaths after inter-hospital transfer” and “deaths after admission,” which showed the highest preventable trauma death rates in a prior study.⁴ We focused on identifying OFIs that can be used as a basis for establishing corrective action plans to reduce preventable trauma death rates. Based on previous studies and guidelines,^{1,10,13-15} OFIs were identified and listed according to the main cause of death: hemorrhage, sepsis/MODS, CNS injury, respiratory failure, and cardiac issues.

Of the patients who experienced a preventable trauma death with hemorrhage as the cause of death, delayed surgery for hemostasis or radiological intervention was the most commonly mentioned factor in relation to “opportunities for improvements,” followed by delayed blood transfusion. Although not presented in the tables, surgery or intervention could not be performed in 21 patients among those with action taken for hemorrhage as the main cause of death; moreover, 17 patients in whom surgery or intervention were performed, there was a median delay of 2 hours and 56 minutes after admission, in providing hemostatic measures. The medical staff failed to perform hemostatic measures within an hour in all cases except for three cases that occurred in an RTC. This result is consistent with that of a prior report, which showed that delayed hemostatic measures were the most common problem found in relation to preventable trauma deaths, even in mature trauma systems.^{11,14,15} Coping with deadly traumatic hemorrhages is a very complex process requiring the involvement of many departments. Many previous reports have shown that the occurrence of preventable trauma deaths can be reduced by establishing effective quality improvement.¹⁶⁻¹⁸ There was a report of a quality improvement case in which the above-mentioned problems were rectified by revising the initial transportation guidelines while transporting severe trauma patients with a potential risk of death from hemorrhage, to a RTC that could perform emergency surgeries or radiological interventions, as well as by educating and training emergency medical care teams with the contents.¹⁹ Issues related to blood transfusions were found in many patients who experienced preventable trauma deaths; in six patients, the medical staff failed to

perform transfusions until the patient's death. Even when performed, it was substantially delayed, with a median time of 64 minutes. This may be due to a lack of provision of emergency transfusions and the need for massive transfusion protocols in facilities other than RTCs.²⁰ The lack of this type of protocol can have a grave impact on treatment outcomes in those with severe traumatic hemorrhages. Therefore, designated trauma centers must have detailed protocols.²¹ A recent study at Penn Medicine Lancaster General Health in 2016 evaluated opportunities for improving the massive transfusion protocol when 11 patients were found to have massive transfusion problems. Therefore, a standard training procedure known as crew resource management (CRM) was implemented, which was associated with decreased mortality.²² The concept of CRM, originally established for the aviation industry quality management attributes most mistakes to the lack of communication and coordination rather than personal error. The CRM approach could be used to solve problems that occur during transfusion processes. In addition, this type of problem solving may need to include a more detailed massive transfusion protocol.^{23,24}

In the case of sepsis/MODS and CNS injury—the second and third most common causes of death after hemorrhage, respectively—delays in treatment at the hospital stage were identified as major OFIs. Regarding cases of sepsis/MODS, the development of corrective action plans should focus on the proper and timely identification and removal of sources of infection that may lead to sepsis. These sources include contamination in the early period that originated from the trauma itself and secondary infections in the late stage that were developed during intensive care. Traumatic brain injury was the main cause of death of all CNS injury patients. In these cases, delay in craniotomy was identified as a major OFI from the review of preventable trauma deaths caused by subdural or epidural brain hemorrhage requiring emergency craniotomy. When craniotomy was performed, the average time until craniotomy after arriving at the final destined hospital was more than double of that recommended by the WHO guidelines for trauma quality improvement programs (“within 4 hours”).¹ In fact, when considering the time spent in the process of transfer before arrival at the final destined hospital, this delay becomes even longer and more serious. Furthermore, 43% (6/14) of CNS injury patients died before craniotomy could be performed. Therefore, reducing the delay in craniotomy is one of the most urgent OFIs identified during the process of early trauma resuscitation that should be addressed to reduce preventable trauma death rates.

This study has some limitations. First, we conducted this study using data from an administrative district of Korea; therefore, the generalization of our results to the entire country would be limited, even though the studied region is one of the largest and most populated provinces in Korea. Second, the retrospective study design precluded the analysis of unrecorded factors or missing values. Third, the evaluation of “preventability” relied entirely on the analysis by expert panels, which has limitations in objective reproducibility, although the Cohen’s Kappa index indicated fair reliability similar to that in previous studies.^{4,25}

The regional preventable trauma death rate in Gyeonggi Province in 2017 was found to be 17.0%. Although this is an improvement from that found in prior studies, many problems were discovered in the inter-hospital transfer stage and hospital stage in emergency medical care facilities other than an RTC. Further, several OFIs regarding bleeding control were identified. To address this, close cooperation among the existing RTCs, other emergency medical care facilities, 119 EMS, and local governments is crucial, and an inclusive regional trauma system needs to be established to enhance further advances in trauma care.

ACKNOWLEDGMENTS

The authors especially thank all the hospitals that participated in this study and the Gyeonggi Province who assisted the research.

SUPPLEMENTARY MATERIAL

Supplementary Data 1

Preventable trauma death pre-review & panel review form

[Click here to view](#)

REFERENCES

1. World Health Organization. *Guidelines for Trauma Quality Improvement Programmes*. Geneva, Switzerland: WHO Press; 2009.
2. Mock C, Joshipura M, Arreola-Risa C, Quansah R. An estimate of the number of lives that could be saved through improvements in trauma care globally. *World J Surg* 2012;36(5):959-63.
[PUBMED](#) | [CROSSREF](#)
3. Cales RH, Trunkey DD. Preventable trauma deaths. A review of trauma care systems development. *JAMA* 1985;254(8):1059-63.
[PUBMED](#) | [CROSSREF](#)
4. Jung K, Kim I, Park SK, Cho H, Park CY, Yun JH, et al. Preventable trauma death rate after establishing a national trauma system in Korea. *J Korean Med Sci* 2019;34(8):e65.
[PUBMED](#) | [CROSSREF](#)
5. Kim Y, Jung KY, Cho KH, Kim H, Ahn HC, Oh SH, et al. Preventable trauma deaths rates and management errors in emergency medical system in Korea. *J Korean Soc Emerg Med* 2006;17(5):385-94.
6. Kim Y. *Assessing Preventable Trauma Death Rate and Improving Trauma Center Administration*. Seoul, Korea: Ministry of Health & Welfare; Research and Development Business Foundations in Seoul National University; 2017.
7. Korean Statistical Information Service. Annual population demographics. http://kosis.kr/statisticsList/statisticsListIndex.do?menuId=M_01_01&vwcd=MT_ZTITLE&parmTabId=M_01_01. Updated 2019. Accessed March 1, 2020.
8. Jung KY, Kim JS, Kim Y. Problems in trauma care and preventable deaths. *J Korean Soc Emerg Med* 2001;12(1):45-56.
9. Motomura T, Mashiko K, Matsumoto H, Motomura A, Iwase H, Oda S, et al. Preventable trauma deaths after traffic accidents in Chiba Prefecture, Japan, 2011: problems and solutions. *J Nippon Med Sch* 2014;81(5):320-7.
[PUBMED](#) | [CROSSREF](#)
10. Sanddal TL, Esposito TJ, Whitney JR, Hartford D, Taillac PP, Mann NC, et al. Analysis of preventable trauma deaths and opportunities for trauma care improvement in Utah. *J Trauma* 2011;70(4):970-7.
[PUBMED](#) | [CROSSREF](#)
11. Teixeira PG, Inaba K, Hadjizacharia P, Brown C, Salim A, Rhee P, et al. Preventable or potentially preventable mortality at a mature trauma center. *J Trauma* 2007;63(6):1338-46.
[PUBMED](#) | [CROSSREF](#)
12. Nirula R, Maier R, Moore E, Sperry J, Gentilello L. Scoop and run to the trauma center or stay and play at the local hospital: hospital transfer's effect on mortality. *J Trauma* 2010;69(3):595-9.
[PUBMED](#) | [CROSSREF](#)
13. Rotondo MF, Cribari C, Smith RS; American College of Surgeons Committee on Trauma. *Resources for Optimal Care of the Injured Patient*. 6th ed. Chicago, IL: American College of Surgeons; 2014.
14. O'Reilly D, Mahendran K, West A, Shirley P, Walsh M, Tai N. Opportunities for improvement in the management of patients who die from haemorrhage after trauma. *Br J Surg* 2013;100(6):749-55.
[PUBMED](#) | [CROSSREF](#)

15. Pfeifer R, Tarkin IS, Rocos B, Pape HC. Patterns of mortality and causes of death in polytrauma patients--has anything changed? *Injury* 2009;40(9):907-11.
[PUBMED](#) | [CROSSREF](#)
16. Davenport RA, Tai N, West A, Bouamra O, Aylwin C, Woodford M, et al. A major trauma centre is a specialty hospital not a hospital of specialties. *Br J Surg* 2010;97(1):109-17.
[PUBMED](#) | [CROSSREF](#)
17. Celso B, Tepas J, Langland-Orban B, Pracht E, Papa L, Lottenberg L, et al. A systematic review and meta-analysis comparing outcome of severely injured patients treated in trauma centers following the establishment of trauma systems. *J Trauma* 2006;60(2):371-8.
[PUBMED](#) | [CROSSREF](#)
18. Liberman M, Mulder DS, Jurkovich GJ, Sampalis JS. The association between trauma system and trauma center components and outcome in a mature regionalized trauma system. *Surgery* 2005;137(6):647-58.
[PUBMED](#) | [CROSSREF](#)
19. Chardbunchachai W, Suppachutikul A, Santikarn C. *Development of Service System for Injury Patients by Utilizing Data from the Trauma Registry*. Khon Kaen, Thailand: Trauma and Critical Care Center, Khon Kaen Regional Hospital; 2002.
20. Hwang K, Kwon J, Cho J, Heo Y, Lee JC, Jung K. Implementation of trauma center and massive transfusion protocol improves outcomes for major trauma patients: a study at a single institution in Korea. *World J Surg* 2018;42(7):2067-75.
[PUBMED](#) | [CROSSREF](#)
21. Malone DL, Hess JR, Fingerhut A. Massive transfusion practices around the globe and a suggestion for a common massive transfusion protocol. *J Trauma* 2006;60(6 Suppl):S91-6.
[PUBMED](#) | [CROSSREF](#)
22. Oriol MD. Crew resource management: applications in healthcare organizations. *J Nurs Adm* 2006;36(9):402-6.
[PUBMED](#) | [CROSSREF](#)
23. Brohi K, Eaglestone S. *Traumatic Coagulopathy and Massive Transfusion: Improving Outcomes and Saving Blood*. Southampton, UK: NIHR Journals Library; 2017.
24. Vernon T, Morgan M, Morrison C. Bad blood: a coagulopathy associated with trauma and massive transfusion review. *Acute Med Surg* 2019;6(3):215-22.
[PUBMED](#) | [CROSSREF](#)
25. MacKenzie EJ, Steinwachs DM, Bone LR, Floccare DJ, Ramzy AI; The Preventable Death Study Group. Inter-rater reliability of preventable death judgments. *J Trauma* 1992;33(2):292-302.
[PUBMED](#) | [CROSSREF](#)