

Analysis of Relationship between Levofloxacin and Corrected QT Prolongation Using a Clinical Data Warehouse

Man Young Park¹, Eun Yeob Kim, MPH², Young Ho Lee, PhD³, Woojae Kim¹, Ku Sang Kim^{1,6}, Seung Soo Sheen, MD⁴, Hong Seok Lim, MD, PhD⁵, Rae Woong Park, MD, PhD¹

¹Department of Biomedical Informatics, Ajou University School of Medicine, Suwon; ²Department of Health Administration, School of Medical Information, Seokang University, Gwangju; ³Department of Information Technology, Gachon University of Medicine and Science, Incheon; Departments of ⁴Pulmonary and Critical Care Medicine, ⁵Cardiology, and ⁶Surgery, Ajou University School of Medicine, Suwon, Korea

Objective: Objective: The aim of this study was to examine whether or not levofloxacin has any relationship with QT prolongation in a real clinical setting by analyzing a clinical data warehouse of data collected from different hospital information systems. **Methods:** Electronic prescription data and medical charts from 3 different hospitals spanning the past 9 years were reviewed, and a clinical data warehouse was constructed. Patients who were both administrated levofloxacin and given electrocardiograms (ECG) were selected. The correlations between various patient characteristics, concomitant drugs, corrected QT (QTc) prolongation, and the interval difference in QTc before and after levofloxacin administration were analyzed. **Results:** A total of 2,176 patients from 3 different hospitals were included in the study. QTc prolongation was found in 364 patients (16.7%). The study revealed that age (OR 1.026, $p < 0.001$), gender (OR 0.676, $p = 0.007$), body temperature (OR 1.267, $p = 0.024$), and cigarette smoking (OR 1.641, $p = 0.022$) were related with QTc prolongation. After adjusting for related factors, 12 drugs concomitant with levofloxacin were associated with QTc prolongation. For patients who took ECGs before and after administration of levofloxacin during their hospitalization ($n = 112$), there was no significant difference in QTc prolongation. **Conclusions:** The age, gender, body temperature, cigarette smoking and various concomitant drugs might be related with QTc prolongation. However, there was no definite causal relationship or interaction between levofloxacin and QTc prolongation. Alternative surveillance methods utilizing the massive accumulation of electronic medical data seem to be essential to adverse drug reaction surveillance in future.

Keywords: Long QT Syndrome, Ofloxacin, Data Mining, Product Surveillance, Post-marketing, Hospital Information Systems

Received for review: January 28, 2011

Accepted for publication: March 25, 2011

Corresponding Author

Rae Woong Park, MD, PhD

Department of Biomedical Informatics, Ajou University School of Medicine, Wonchon-dong, Yeongtong-gu, Suwon 442-749, Korea.
Tel: +82-31-219-5342, Fax: +82-31-219-4472, E-mail: veritas@ajou.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

© 2011 The Korean Society of Medical Informatics

1. Introduction

QT interval on the surface electrocardiogram (ECG) is the period from depolarization to repolarization of ventricle, and QT interval can be calculated in the ECG by measuring the distance between the start of the Q wave and the end of the T wave. Prolongation of the QT interval is known to possibly cause ventricular arrhythmia, including tachycardia, fibrillation, and torsades de pointes (TdP), resulting in syncope or sudden death [1-3]. Because of fatal outcome, the possibility of QT prolongation has been a very important issue. Many non-cardiovascular drugs are suspected

to cause QT prolongation. They include some antidepressants (amitriptyline, fluoxetine, doxepine, venlafaxine, etc.), antimicrobials (azithromycin, levofloxacin, roxithromycin, etc.), antihistamines (astemizole, clemastine, etc.), antipsychotics (clozapine, quetiapine, sultopride, etc.) and so on [1, 4-6]. There have been label changes or market withdrawal of drugs found to carry high risk of QT prolongation [3,7], and these drugs include terfenadine, astemizole, grepafloxacin, terodiline, droperidol, lidoflazine, sertindole, levomethadyl, and cisapride.

The reports of adverse drug reaction (ADR) during pre-marketing clinical trials are very limited, because of limitation of the number of subjects, clinical setting and age groups. For example, phase 3 clinical trials are usually performed with only 1,000 to 3,000 subjects, and children, old and pregnant women are usually excluded from the studies [8]. Therefore, the monitoring of safety of drugs after marketing is mandatory to find any harmful reactions that were not observed prior to approval. The safety of drugs after marketing is continuously monitored through epidemiologic studies, such as spontaneous report, chart review, cohort study and case report [9]. Some drugs, such as cisapride, celecoxib and rofecoxib were withdrawn from the market due to fatal ADRs, even though they showed no noticeable ADRs in clinical trials [10].

Recently, the electronic medical record (EMR) and computerized physician order entry (CPOE) system adoption rate has been ever increasing [11,12]. As a result, there are many efforts to use EMR and CPOE data for the capture of ADR [13]. Use of data warehouse in clinical areas has also been reported [14,15]. Data warehouse (DW) is a database repository for the analytical purposes from the operational database. Szirbik et al. [16] suggests a general method for constructing a clinical DW. Sheen et al. [17] reported that overdose rate in patients with moderate to severe renal insufficiency was 28.2%, when tertiary teaching hospital's 4 years prescription data in a DW were analyzed. Zhang et al. [15] analyzed clinical prescription data in a DW and reported that concomitant administration of aminoglycoside and cephalosporin increased nephrotoxicity, and Hinrichsen et al. [18] constructed an electronic vaccine adverse reaction reporting system that was interfaced to the EMR system, and found 6-fold increase in vaccine adverse reaction reports compared to traditional self reporting method.

Levofloxacin, an antibiotic of the fluoroquinolone drug class had been reported to cause QT prolongation in animal studies [19]. There have been several reports that levofloxacin is associated with corrected QT (QTc) prolongation [20-22]. However, some study has reported that there is no effect

of levofloxacin on QTc prolongation [23]. Therefore, the effect of levofloxacin on QTc prolongation remains still controversial.

This study aims to examine whether levofloxacin has any relation with QT prolongation in real clinical setting by using a clinical DW, which contained past 9 years electronic prescription data from 3 different general hospitals. We analyzed the relationship between epidemiologic and clinical characteristics of the subjects with QT prolongation. The relationship between the concomitant drugs with levofloxacin and QT prolongation was analyzed, after adjusting epidemiologic and clinical characteristics of the subjects. Finally, the difference in QT prolongation between pre- and post-administration of levofloxacin was analyzed.

II. Methods

1. Data Preparation and Study Population

Three hospitals joined in this study. A hospital is a 1,089-bed tertiary teaching hospital, B hospital is a 1,198-bed tertiary teaching hospital, and C hospital is a 131-bed general hospital. The study period ranged 9 years for A hospital (1998-2006), and 5 years for B and C hospitals (2002-2006). This study was reviewed and approved by the Institutional review board committees.

The method how to make data mart for this study was partly introduced in the previous work in detail [24]. Briefly, patients who were administrated subject drug (levofloxacin) during their hospitalization within the period mentioned above were selected. Then, every prescription and ECG examination data on patients who were selected was transformed and loaded into a data base (MS-SQL 2000, Microsoft, USA). The resulting data included 136,146 levofloxacin prescriptions for 21,950 patients. Of patients who were administrated levofloxacin and whose ECG was taken during the hospitalization, only patients whose age was 18 years or older were selected ($n = 3,854$) (Figure 1).

Patients' epidemiologic and clinical data were collected by manually reviewing the available medical charts ($n = 2,903$). The epidemiologic and clinical data included age, gender, height, body weight, blood pressure, body temperature, heart rate, body mass index (BMI), cigarette smoking, alcohol drinking, presence of chronic disease (hypertension, diabetes, hepatitis, and tuberculosis), final discharge diagnoses, operation and treatment procedures. ECG results were also reviewed manually. By reviewing the discharge summary, one main discharge diagnosis and associate diagnoses up to 10, surgical and treatment procedures up to 7 were examined. De-identification of patients' personal data was per-




Figure 1. Case selection process and scope of the study. A data warehouse, which contains electronic prescription data from 3 different general hospitals, were constructed. Total of 2,176 patients were selected for the analyses. QTc: corrected QT interval.

formed prior to analysis.

For the QTc measurement of hospital A, it was automatically calculated by the ECG device using Bazett's formula [25]. Because the QTc was not available from the ECG device, the QT value was manually measured and then QTc was calculated using Bazett's formula for 2 other hospitals (B, C hospital).

Patients with disease that directly affects ECG wave and QT, such as ischemic heart disease, heart failure, myocarditis, mitral valve prolapsed, atrial fibrillation, cerebrovascular disease, rheumatic fever, anorexia nervosa, starvation, hypothyroidism, hypomagnesemia or renal failure, were excluded from the study (n = 645). Patients with unreliable QTc in ECG due to base-line swing, base-line noise, T-inversion, flat T wave, U-wave, irregular RR interval, premature ventricular contracture due to myocarditis, myocardial infarction, heart failure, pericarditis, atrial fibrillation and coronary arteriosclerosis were also excluded (n = 82). The resulting total number of subjects for the study was 2,176 (M = 1,316, F = 860).

Of 2,176 subjects, 588 patients were selected for the analysis of effects of other concomitant drugs with levofloxacin on QTc prolongation. Concomitant drugs that were 50 times or

more frequently prescribed were selected. Multiple logistic regression for each drug was performed after adjusting the effect of epidemiologic and clinical variables.

On the case-series study design, difference in QTc prolongation before and after levofloxacin administration was analyzed. Patients who had both records on ECG examination before and after levofloxacin were selected (n = 112). It was regarded as 'pre-levofloxacin administration' when a patient was not administrated levofloxacin more than 2 days before ECG examination. On the other hand, it was considered as 'post-levofloxacin administration' when levofloxacin administrated and ECG examination occurred on the same day.

2. Definition of QTc Prolongation

For QTc interval prolongation standards in terms of gender classification, the QTc interval prolongation for men was set at QTc > 450 msec (normal 430 msec + borderline 20 msec), and QTc > 470 msec for women (normal 450 msec + borderline 20 msec) [26].

3. Statistical Analysis

Mann-Whitney and chi-square test were performed to examine the difference in QTc prolongation by clinical or

epidemiologic characteristics of patients. Then multiple logistic regression was performed with the previous major factors affecting QTc prolongation. To reveal the effects of other concomitant drugs with levofloxacin on QTc, multiple logistic regression was performed after adjusting major clinical and epidemiologic factors. Wilcoxon signed rank test was performed in order to determine whether there was difference in QTc prolongation before and after levofloxacin administration. SPSS ver. 12.0 (SPSS Inc., Chicago, IL, USA) was used for the analyses.

III. Results

The analysis showed mean age of 52.5 years (52.7 ± 17.1 for men, 52.3 ± 17.9 for women), mean height of 163.7 cm (168.7 ± 6.5 for men, 156.0 ± 6.7 for women), mean body weight of 60.8 kg (64.5 ± 11.2 for men, 55.0 ± 10.0 for women), mean BMI of 22.6 kg/m² (22.6 ± 3.4 for men, 22.6 ± 3.8 for women), mean systolic blood pressure of 119.2 mmHg (120.0 ± 16.8 for men, 117.9 ± 18.1 for women), mean diastolic blood pressure of 74.5 mmHg (75.0 ± 11.4 for men, 73.7 ± 11.1 for women), mean body temperature of 36.7°C (36.7 ± 0.62 for men, 36.7 ± 0.64 for women), mean heart rate of 80.9 beats/min (80.2 ± 19.2 for men, 81.9 ± 17.5 for women), and mean QTc interval of 423.3 msec (420.7 ± 39.3 for men, 427.2 ± 39.0 for women) (Table 1).

1. Association of Patients' Characteristics with QTc

QTc prolongation was found in 256 (19.4%) out of 1,316 for

men and 108 (12.6%) out of 860 for women who had been administrated levofloxacin during their hospitalization. There was difference in QTc prolongation by age for men and women ($p < 0.001$), height for women ($p < 0.001$), body weight for men ($p = 0.001$), BMI for men ($p = 0.018$), systolic blood pressure ($p = 0.013$), body temperature for men ($p = 0.009$), heart rate for both ($p < 0.001$), cigarette smoking for men ($p = 0.016$) and alcohol drinking for men ($p = 0.057$) at a significance level of 0.1 (Table 2).

With these variables, forward stepwise multiple logistic regression analysis revealed that age ($p < 0.001$), gender ($p = 0.007$), body temperature ($p = 0.024$) and cigarette smoking ($p = 0.022$) were associated with QTc prolongation (Table 3).

2. Association of Concomitant Drugs with QTc

Of 720 drugs jointly administered with levofloxacin, 32 drugs that were prescribed 50 times or more frequently were selected (Table 4).

Chi-square test revealed that 18 drugs were associated with QTc prolongation at the significance level of 0.1. The logistic regression revealed that the following drugs were associated with QTc prolongation, even after adjusting the effect of major variables (age, gender, body temperature and cigarette smoking): furosemide (odd ratio [OR], 5.8; $p < 0.001$), magnesium sulfate (OR, 7.8; $p < 0.001$), propacetamol (OR, 4.1; $p < 0.001$), sucralfate (OR, 2.5; $p = 0.019$), aminophylline (OR, 2.6; $p = 0.004$), vecuronium bromide (OR, 6.0; $p < 0.001$), heparin sodium (OR, 3.7; $p = 0.014$), human serum albumin (OR, 3.8; $p < 0.001$), isosorbide dinitrate (OR, 4.6; $p = 0.003$), morphine (OR, 4.4; $p < 0.001$), clindamycin (OR, 4.9; $p < 0.001$), and amikacin sulfate (OR, 3.0; $p = 0.004$) (Table 5).

3. Difference in QTc after Levofloxacin Administration

Patients who took ECGs before and after administration of levofloxacin during their hospitalization were selected from the clinical data mart. The Wilcoxon signed rank test was performed on the patients extracted to determine whether there was QTc interval differences between pre and post levofloxacin administration (Table 6).

There were total 112 pre and post comparable patients (consisting of 76 males [67.9%], and 36 females [32.1%]). The pre levofloxacin administration QTc mean was 432.9 msec (± 36.8), while the post QTc mean was 434.5 msec (± 33.3). Seventy-seven patients (68.8%) were 'Normal to Normal,' 12 patients (10.7%) were 'Normal to Prolonged,' 10 patients (11.6%) were 'Prolonged to Normal,' and 13 patients (11.6%) were 'Prolonged to Prolonged.' Wilcoxon signed rank test revealed that there was no statistically significant QTc prolongation difference between pre and post levofloxa-

Table 1. Characteristics of the study population (n = 2,176)

	Male (n = 1,316)	Female (n = 860)	p-value
Age	52.7 ± 17.1	52.3 ± 17.9	0.563
Height (cm)	168.7 ± 6.5	156.0 ± 6.7	<0.001
Body weight (kg)	64.5 ± 11.2	55.0 ± 10.0	<0.001
Body mass index (kg/m ²)	22.6 ± 3.4	22.6 ± 3.8	0.896
Systolic blood pressure (mmHg)	120.0 ± 16.8	117.9 ± 18.1	0.006
Diastolic blood pressure (mmHg)	75.0 ± 11.4	73.7 ± 11.1	0.006
Body temperature (°C)	36.7 ± 0.62	36.7 ± 0.64	0.789
Heart rate (beats/min)	80.2 ± 19.2	81.9 ± 17.5	0.330
QTc interval (msec)	420.7 ± 39.3	427.2 ± 39.0	<0.001

Values are presented as mean ± SD.

QTc: corrected QT interval, SD: standard deviation.

Table 2. Comparison of study populations' characteristics by QTc

Continuous variables	QTc for men ^a		p-value ^b	QTc for women ^a		p-value ^b
	Normal (n = 1,060)	Prolonged (n = 256)		Normal (n = 752)	Prolonged (n = 108)	
Age (yr)	51.4 ± 17.1	58.3 ± 16.2	<0.001	50.8 ± 17.9	62.3 ± 14.6	<0.001
Height (m)	168.9 ± 6.5	167.9 ± 6.8	0.111	156.3 ± 6.6	153.6 ± 6.6	<0.001
Body weight (kg)	65.1 ± 11.2	62.0 ± 10.6	<0.001	55.2 ± 10.1	53.2 ± 9.3	0.118
Body mass index (kg/m ²)	22.7 ± 3.4	22.1 ± 3.3	0.018	22.6 ± 3.9	22.5 ± 3.5	0.904
Systolic pressure (mmHg)	119.7 ± 16.0	121.5 ± 19.8	0.388	117.2 ± 17.6	122.7 ± 20.5	0.100
Diastolic pressure (mmHg)	75.2 ± 11.2	74.5 ± 12.2	0.302	73.6 ± 11.1	74.1 ± 10.9	0.348
Body temperature (°C)	36.7 ± 0.6	36.7 ± 0.7	0.009	36.7 ± 0.6	36.7 ± 0.7	0.586
Heart rate (beats/min)	77.7 ± 17.9	91.0 ± 20.5	<0.001	80.8 ± 17.1	89.7 ± 18.7	<0.001
QTc interval (msec)	406.6 ± 26.4	479.1 ± 28.9	<0.001	417.1 ± 28.5	498.0 ± 27.7	<0.001

Categorical variables		No. of QTc for men		p-value ^c	No. of QTc for women		p-value ^c
		Normal	Prolonged		Normal	Prolonged	
Cigarette smoking	No	951	216	0.016	718	102	0.413
	Yes	100	38		25	5	
Alcohol drinking	No	972	225	0.057	726	105	1.000
	Yes	80	29		17	2	
Diabetes	No	962	227	0.327	669	93	0.311
	Yes	90	27		74	14	
Hypertension	No	940	221	0.316	651	90	0.352
	Yes	112	33		92	17	
Tuberculosis	No	946	220	0.141	662	90	0.142
	Yes	106	34		80	17	
Hepatitis	No	1,041	253	0.480	731	104	0.420
	Yes	11	1		12	3	

QTc: corrected QT interval, SD: standard deviation.

^aValues are presented as mean ± SD. ^bMann-Whitney test. ^cChi-square test.

Table 3. Association of factors with QTc prolongation

	OR	95% CI	p-value
Age (yr)	1.026	1.017-1.034	<0.001
Gender (female)	0.676	0.509-0.899	0.007
Body temperature (°C)	1.267	1.031-1.555	0.024
Cigarette smoking	1.641	1.076-2.503	0.022

OR: odds ratio, CI: confidence interval.

cin administration ($p = 0.353$).

IV. Discussion

1. Main Findings

The study revealed that age, gender, body temperature and

cigarette smoking are related with QTc interval prolongation. This result is consistent with previous studies indicating that QTc is related with age, gender, body temperature [27] and cigarette smoking [28].

Even after adjusting the major factors, 12 drug types still showed significant correlation. However, because of the limitation of study design, it is not clear whether such correlations are caused by any interaction, or by associated conditions or by concomitant drugs itself. Furosemide and clindamycin have already been known to be related with QTc prolongation [3]. Isosorbide dinitrate is usually used in angina pectoris, which is related with QT interval change [29]. Morphine is a strong pain-killer used to relieve pain after surgery or cancer pain, and it is likely that it has a correlation to QTc prolongation in diverse situations where it must be administered. Vecuronium bromide is a drug used

Table 4. Association of concomitant drugs with QTc prolongation in combination with levofloxacin

Drug	Drug administration	No. of QTc		p-value ^a
		Normal	Prolonged	
Tramadol	-	353	79	0.753
	+	125	31	
Furosemide	-	440	78	<0.001
	+	38	32	
Acetylcysteine	-	389	72	<0.001
	+	89	38	
Ketorolac tromethamine	-	374	87	0.947
	+	104	23	
Cimetidine	-	408	87	0.139
	+	70	23	
Pethidine	-	383	92	0.479
	+	95	18	
Ranitidine	-	438	92	0.018
	+	40	18	
Pancreatine	-	407	94	1.000
	+	71	16	
Metoclopramide	-	412	95	1.000
	+	66	15	
Magnesium sulfate	-	459	86	<0.001
	+	19	24	
Propacetamol	-	440	86	<0.001
	+	38	24	
Bromhexine	-	443	90	0.001
	+	35	20	
Sucralfate	-	457	96	0.002
	+	21	14	
Aminophylline	-	448	86	<0.001
	+	30	24	
Vecuronium bromide	-	460	85	<0.001
	+	18	25	
Heparin sodium	-	466	102	0.028
	+	12	8	
Human serum albumin	-	455	89	<0.001
	+	23	21	
Tranexamic acid	-	428	103	0.258
	+	50	7	
Isosorbide dinitrate	-	470	99	<0.001
	+	8	11	

Table 4. Continued

Drug	Drug administration	No. of QTc		p-value ^a
		Normal	Prolonged	
Morphine	-	457	88	<0.001
	+	21	22	
Isoniazide	-	423	101	0.401
	+	55	9	
Clindamycin	-	453	87	<0.001
	+	25	23	
Codeine phosphate	-	438	99	0.719
	+	40	11	
Itraconazole	-	423	99	0.777
	+	55	11	
Digoxin	-	475	105	0.006
	+	3	5	
Tiropramide	-	453	106	0.651
	+	25	4	
Roxatidine acetate	-	464	103	0.143
	+	14	7	
Amlodipine besylate	-	462	100	0.017
	+	16	10	
Atrophine sulfate	-	477	110	1.000
	+	1	0	
Amikacin sulfate	-	453	97	0.02
	+	25	13	
Caroverine	-	429	105	0.092
	+	49	5	
Prednisolone	-	455	101	0.241
	+	23	9	

QTc: corrected QT interval. ^aChi-square test.

to relax muscles during operations. According to the study by Radhakrishnan et al. [30] there has been cases of QT interval prolongation in patients due to heparin-induced hypocalcaemia. Magnesium sulfate is known to be effective in the treatment of ventricular arrhythmias. However, there still lack reports of correlation between QT prolongation and aminophylline, propacetamol, sucralfate, and amikacin sulfate. Individual drug itself or joint administration of different drugs may be related with QTc prolongation or the clinical

Table 5. Association of concomitant drugs with QTc prolongation in combination with levofloxacin, after adjusting the major variables: age, gender, body temperature and cigarette smoking

Drug	OR	95% CI	p-value
Furosemide	5.8	3.2-10.4	<0.001
Magnesium sulfate	7.8	3.7-16.4	<0.001
Propacetamol	4.1	2-7.6	<0.001
Sucralfate	2.5	1.2-5.3	0.019
Aminophylline	2.6	1.4-4.9	0.004
Vecuronium bromide	6.0	3.0-11.9	<0.001
Heparin sodium	3.7	1.3-10.7	0.014
Human serum albumin	3.8	1.9-7.6	<0.001
Isosorbide dinitrate	4.6	1.7-12.6	0.003
Morphine	4.4	2.2-8.8	<0.001
Clindamycin	4.9	2.5-9.5	<0.001
Amikacin sulfate	3.0	1.4-6.5	0.004

OR: odds ratio, CI: confidence interval.

condition in which the relevant drugs administered can also be related with the QTc prolongation. Nykamp et al. [20] has studied on QTc prolongation associated with levofloxacin. He concluded that the use of levofloxacin alone may cause QT interval prolongation. Further study is needed for more reasonable explanation.

One of the essential points of this study was to develop and discover research methods utilizing heavily accumulating medical data in electronic form. EMR and CPOE systems are fast adopted by many hospitals [11,12]. However, the use of data warehouse or data mart in the field of post-market surveillance (PMS) is in its early stage. At present, much of actual clinical electronic databases are created within the hospitals, nevertheless, most of the clinically important patient information are records on paper or non-standardized data on the electronic database. Consequently, therefore, the information had to be obtained manually. Integrating electronic medical data from different hospitals is a difficult, tedious and multistep process not only in terms of technical problems of integrating heterogeneous database schema, but also problems of institutional policies, privacy and security problems. The detailed processes of data warehouse or data mart construction have been reported elsewhere [24]. Although massive data-driven analysis, such as the present study, has not been described to explain a causal relationship compared to well-designed clinical trial, nonetheless, it can be useful to get a simple and quick insight into a problem.

Table 6. Comparison of QTc between before and after levofloxacin administration

QTc (before → after)	Male	Female	Total (%)	
Normal → Normal	51	26	77 (68.8)	
Normal → Prolonged	8	4	12 (10.7)	
Prolonged → Normal	5	5	10 (8.9)	
Prolonged → Prolonged	12	1	13 (11.6)	
Total	76	36	112 (100.0)	
Mean QTc (msec)	Before	428.8 (± 30.1)	441.6 (± 47.3)	432.9 (± 36.8)
	After	434.7 (± 35.3)	434.0 (± 29.2)	434.5 (± 33.3)
	p-value ^a	0.186	0.802	0.353

QTc: corrected QT interval. ^aWilcoxon signed rank test.

2. Limitations

The present results carry several limitations. As a retrospective study on general in-patients, major confounding factors could not be pre-adjusted, and details of diseases for hospital admission, base disease, major history, various drug administration and treatments while in hospital are heterogeneous. Taking these limitations into account, the major clinical indicators of subjects (age, gender, body temperature, cigarette smoking) were adjusted during the analysis process, but limitations may still be inherent. As a study just on the exposed group to levofloxacin without a control group, the relationship between cause and result could not be examined. As a retrospective study, this research was dependent mainly on the records in relation to the measurement of major variables, therefore adequate management of measurement accuracy and precision could not be evaluated. Moreover, systematic measurement errors of the participating institution could also exist. For example, there are many types of ECG devices which would be different for each department and hospital. Because the types of ECG devices are different between hospitals and some may even be using non-automated ECG devices, there could be inter-observer and intra-observer variability when manually measuring QT interval. Because the conditions of patients were so diverse, and representative diagnosis and ECG values were selected, thus giving rise to some variation in the information.

In this study, subjects who had heart diseases according to the main-diagnosis and sub-diagnosis records were excluded from the study population. However, drugs for cardiovascular diseases were found in the concomitant drugs. It seems that the discharge diagnoses of patients may not in some

cases be representative for the present illness.

There was no statistically significant difference in QTc prolongation between before and after administration of levofloxacin. However, significance may not be clear in this massive data-driven study, because of heterogeneous nature of the data set obtained from real clinical environments. For definite analysis, a more sophisticated clinical data mart for PMS would be helpful. We are in a process to construct a more well-designed clinical data mart for PMS.

Although we performed this study to identify possible association between levofloxacin and QTc prolongation, a lot of time and extensive labor was consumed due to required many manual processes. A well designed database-driven approach, analyzing the drug-ADR of EMR databases such as comparison of exposed and non-exposed group for a drug with automated process which not need any intervention of specialist, seems essential to identify ADR signals quickly. We are preparing an automated ADR signal detection algorithm and system.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2010-0023402).

References

1. Yap YG, Camm AJ. Drug induced QT prolongation and torsades de pointes. *Heart* 2003; 89: 1363-1372.
2. Moss AJ, Schwartz PJ, Crampton RS, Tzivoni D, Locati EH, MacCluer J, Hall WJ, Weitkamp L, Vincent GM, Garson A Jr. The long QT syndrome: prospective longitudinal study of 328 families. *Circulation* 1991; 84: 1136-1144.
3. Fermini B, Fossa AA. The impact of drug-induced QT interval prolongation on drug discovery and development. *Nat Rev Drug Discov* 2003; 2: 439-447.
4. Woosley RL. Cardiac actions of antihistamines. *Annu Rev Pharmacol Toxicol* 1996; 36: 233-252.
5. Yap YG, Camm AJ. The current cardiac safety situation with antihistamines. *Clin Exp Allergy* 1999; 29 Suppl 1: 15-24.
6. Arizona Center for Education and Research on Therapeutics. QT drug lists by risk groups. Arizona Center for Education and Research on Therapeutics: [cited 2011 Mar 29]. Available from: <http://www.azcert.org/medical-pros/drug-lists/drug-lists.cfm>.
7. Brown AM. Drugs, hERG and sudden death. *Cell Calcium* 2004; 35: 543-547.
8. Ahmad SR. Adverse drug event monitoring at the food and drug administration. *J Gen Intern Med* 2003; 18: 57-60.
9. Waller PC, Evans SJ. A model for the future conduct of pharmacovigilance. *Pharmacoepidemiol Drug Saf* 2003; 12: 17-29.
10. Weaver J, Bonnel RA, Karwoski CB, Brinker AD, Beitz J. GI events leading to death in association with celecoxib and rofecoxib. *Am J Gastroenterol* 2001; 96: 3449-3450.
11. DesRoches CM, Campbell EG, Rao SR, Donelan K, Ferris TG, Jha A, Kaushal R, Levy DE, Rosenbaum S, Shields AE, Blumenthal D. Electronic health records in ambulatory care: a national survey of physicians. *N Engl J Med* 2008; 359: 50-60.
12. Park RW, Shin SS, Choi YI, Ahn JO, Hwang SC. Computerized physician order entry and electronic medical record systems in Korean teaching and general hospitals: results of a 2004 survey. *J Am Med Inform Assoc* 2005; 12: 642-647.
13. Hauben M, Patadia V, Gerrits C, Walsh L, Reich L. Data mining in pharmacovigilance: the need for a balanced perspective. *Drug Saf* 2005; 28: 835-842.
14. Dewitt JG, Hampton PM. Development of a data warehouse at an academic health system: knowing a place for the first time. *Acad Med* 2005; 80: 1019-1025.
15. Zhang Q, Matsumura Y, Teratani T, Yoshimoto S, Mineo T, Nakagawa K, Nagahama M, Kuwata S, Takeda H. The application of an institutional clinical data warehouse to the assessment of adverse drug reactions (ADRs): evaluation of aminoglycoside and cephalosporin associated nephrotoxicity. *Methods Inf Med* 2007; 46: 516-522.
16. Szirbik NB, Pelletier C, Chausalet T. Six methodological steps to build medical data warehouses for research. *Int J Med Inform* 2006; 75: 683-691.
17. Sheen SS, Choi JE, Park RW, Kim EY, Lee YH, Kang UG. Overdose rate of drugs requiring renal dose adjustment: data analysis of 4 years prescriptions at a tertiary teaching hospital. *J Gen Intern Med* 2008; 23: 423-428.
18. Hinrichsen VL, Kruskal B, O'Brien MA, Lieu TA, Platt R. Using electronic medical records to enhance detection and reporting of vaccine adverse events. *J Am Med*

- Inform Assoc 2007; 14: 731-735.
19. Akita M, Shibazaki Y, Izumi M, Hiratsuka K, Sakai T, Kurosawa T, Shindo Y. Comparative assessment of prurifloxacin, sparfloxacin, gatifloxacin and levofloxacin in the rabbit model of proarrhythmia. *J Toxicol Sci* 2004; 29: 63-71.
 20. Nykamp DL, Blackmon CL, Schmidt PE, Roberson AG. QTc prolongation associated with combination therapy of levofloxacin, imipramine, and fluoxetine. *Ann Pharmacother* 2005; 39: 543-546.
 21. Gandhi PJ, Menezes PA, Vu HT, Rivera AL, Ramaswamy K. Fluconazole- and levofloxacin-induced torsades de pointes in an intensive care unit patient. *Am J Health Syst Pharm* 2003; 60: 2479-2483.
 22. Taubel J, Naseem A, Harada T, Wang D, Arezina R, Lorch U, Camm AJ. Levofloxacin can be used effectively as a positive control in thorough QT/QTc studies in healthy volunteers. *Br J Clin Pharmacol* 2010; 69: 391-400.
 23. Tsikouris JP, Peeters MJ, Cox CD, Meyerrose GE, Seifert CF. Effects of three fluoroquinolones on QT analysis after standard treatment courses. *Ann Noninvasive Electrocardiol* 2006; 11: 52-56.
 24. Park MY, Lee YH, Kim EY, Kim WJ, Kam HJ, Choi JP, Han TH, Kang UG, Park RW. A data warehouse based retrospective post-marketing surveillance method: a feasibility test with fluoxetine. *J Korean Soc Med Inform* 2009; 15: 191-199.
 25. Bazett HC. The time relations of the blood-pressure changes after excision of the adrenal glands, with some observations on blood volume changes. *J Physiol* 1920; 53: 320-339.
 26. Viskin S. Long QT syndromes and torsade de pointes. *Lancet* 1999; 354: 1625-1633.
 27. Mattu A, Brady WJ, Perron AD. Electrocardiographic manifestations of hypothermia. *Am J Emerg Med* 2002; 20: 314-326.
 28. Singh K. Effect of smoking on QT interval, QT dispersion and rate pressure product. *Indian Heart J* 2004; 56: 140-142.
 29. Suzuki M, Nishizaki M, Arita M, Ashikaga T, Yamawake N, Kakuta T, Numano F, Hiraoka M. Increased QT dispersion in patients with vasospastic angina. *Circulation* 1998; 98: 435-440.
 30. Radhakrishnan M, Agarwal S, Bithal PK, Gupta V. Heparin-induced transient prolongation of the QT interval during endovascular embolisation of intracranial aneurysm. *J Clin Neurosci* 2006; 13: 489-492.